

CORONARY ARTERY DISEASE

Acute coronary syndromes (NSTEMI and STEMI)

Relevant Guidelines

ACC/AHA 2025 guidelines on ACS

2025 ACC/AHA/ACEP/NAEMSP/SCAI Guideline for the Management of Patients With Acute Coronary Syndromes. *J Am Coll Cardiol.* 2025

AHA/ACC 2021 guidelines on Chest Pain

2021 AHA/ACC/ASE/CHEST/SAEM/SCCT/SCMR Guideline for the Evaluation and Diagnosis of Chest Pain. *Circulation.* 2021

ESC 2023 guidelines on acute coronary syndromes

2023 ESC Guidelines for the management of acute coronary syndromes. *Eur Heart J.* 2023.

ESC/EACTS 2018 guidelines on revascularization

2018 ESC/EACTS Guidelines on myocardial revascularization. *Eur Heart J.* 2018.

ESC 2021 guidelines on heart failure

2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J*

ESC 2021 guidelines on pacing and cardiac resynchronization

2021 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy. *Eur Heart J.* 2021.

ACC/AHA/ACCP/HRS 2023 guideline on AF

2023 ACC/AHA/ACCP/HRS Guideline for the Diagnosis and Management of Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation.* 2023

ACCF/AHA/HRS 2018 guideline on bradycardia and conduction delay

2018 ACC/AHA/HRS Guideline on the Evaluation and Management of Patients With Bradycardia and Cardiac Conduction Delay

AHA/ACC 2026 Guideline on Cardiovascular-Kidney-Metabolic Syndrome

2026 AHA/ACC/ADA/ASN Guideline for the Prevention, Detection, Evaluation, and Management of Cardiovascular-Kidney-Metabolic Syndrome: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol.* 2026;87:e1889-e2007.

ESC 2026 GL on chronic kidney disease

2026 ESC Guidelines for the management of cardiovascular disease and chronic kidney disease, in collaboration with the European Renal Association (ERA). *Eur Heart J.* 2026.

ESC 2024 guidelines on AF

2024 ESC Guidelines for the management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J.* 2024.

ESC 2022 Guidelines on ventricular arrhythmias and prevention of sudden cardiac death

2022 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death. *Eur Heart J.* 2022.

ACC/AHA 2019 Guideline on the Primary Prevention of Cardiovascular Disease

2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease. *J Am Coll Cardiol.* 2019;74:1376-1414.

ESC 2021 Guidelines on cardiovascular disease prevention

2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J.* 2021

ACC/AHA 2026 Guideline on Management of Dyslipidemia

ACC/AHA/AACVPR/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Dyslipidemia: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2026.

ESC 2023 Guidelines in diabetes

2023 ESC Guidelines for the management of cardiovascular disease in patients with diabetes. *Eur Heart J*. 2023.

ESC 2019 Guidelines on Dyslipidaemias

2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J*. 2019.

ESC 2025 Update of Guidelines on Dyslipidaemias

2025 Focused Update of the 2019 ESC/EAS Guidelines for the management of dyslipidaemias. *Eur Heart J*. 2025

ESC 2020 Guidelines on sports

2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease. *Eur Heart J*. 2020.

ESC 2022 Guidelines on cardio-oncology

2022 ESC Guidelines on cardio-oncology developed in collaboration with the European Hematology Association (EHA), the European Society for Therapeutic Radiology and Oncology (ESTRO) and the International Cardio-Oncology Society (IC-OS). *Eur Heart J*. 2022.

ESC 2026 Guidelines on Rehabilitation

2026 ESC Guidelines on cardiac rehabilitation. *Eur Heart J*. 2026.

ESC 2025 Guidelines on pregnancy.

ESC 2025 Guidelines on the management of cardiovascular diseases and pregnancy. *Eur Heart J*. 2025.

Relevant Scientific Statements and Important Position Papers**ESC/ACC/AHA/WHF 2018 Fourth universal definition of myocardial infarction**

Thygesen K, et al. Fourth universal definition of myocardial infarction (2018). *Eur Heart J*. 2018.

AHA 2016 Scientific Statement on STEMI in women

Acute Myocardial Infarction in Women: A Scientific Statement From the American Heart Association. *Circulation*. 2016;133:916-947

AHA 2018 Statement on Spontaneous Coronary Artery Dissection

Spontaneous Coronary Artery Dissection: Current State of the Science: A Scientific Statement From the American Heart Association. *Circulation*. 2018.

AHA 2021 Statement on Cardiogenic Shock

Invasive Management of Acute Myocardial Infarction Complicated by Cardiogenic Shock: A Scientific Statement From the American Heart Association. *Circulation*. 2021;143:e815-e829.

AHA 2021 Statement on Practices in the Cardiac Catheterization Laboratory

Evidence-Based Practices in the Cardiac Catheterization Laboratory: A Scientific Statement From the American Heart Association. *Circulation*. 2021:CIR0000000000000996.

AHA 2021 Statement on Mechanical Complications of Acute MI

Mechanical Complications of Acute Myocardial Infarction: A Scientific Statement From the American Heart Association. *Circulation*. 2021:CIR0000000000000985.

AHA 2022 Scientific Statement on ACS in the Elderly

Management of Acute Coronary Syndrome in the Older Adult Population: A Scientific Statement From the American Heart Association. *Circulation*. 2022

AHA 2023 Scientific Statement on Postoperative Ischaemia after CABG

Considerations on the Management of Acute Postoperative Ischaemia After Cardiac Surgery: A Scientific Statement From the American Heart Association. *Circulation*. 2023;148:442-454.

AHA 2024 Scientific Statement on CYP2C19 Genetic Testing

CYP2C19 Genetic Testing for Oral P2Y12 Inhibitor Therapy: A Scientific Statement From the American Heart Association. *Circulation*. 2024;150:e129-e150.

ESC groups 2026 Statement on sex differences in IHD.

Sex and gender differences in coronary pathophysiology and ischaemic heart disease. *Eur Heart J*. 2026;47:2727-2749.

ESC groups 2026 Statement on antithrombotic therapy

Antithrombotic therapy for secondary prevention in patients with acute coronary syndromes treated with percutaneous coronary intervention: options for personalization to reduce bleeding or ischaemic risks. A Clinical Consensus Statement of the ESC Working Group on Thrombosis, the Association for Acute CardioVascular Care of the ESC, the European Association of Percutaneous Cardiovascular Interventions of the ESC, and the ESC Working Group on Cardiovascular Pharmacology. *EuroIntervention*. 2026;22:e653-e677.

ACC 2026 Report on Antiplatelet Therapy

Antiplatelet Therapy in the Management of Atherosclerotic Cardiovascular Disease: 2026 ACC Scientific Statement: A Report of the American College of Cardiology. *J Am Coll Cardiol*. 2026.

Stable CAD

Relevant Guidelines

ESC 2024 guidelines on chronic coronary syndromes

2024 ESC Guidelines for the diagnosis and management of chronic coronary syndromes. *Eur Heart J.* 2024.

AHA/ACC 2023 guideline on Chronic Coronary Disease

2023 AHA/ACC/ACCP/ASPC/NLA/PCNA Guideline for the Management of Patients With Chronic Coronary Disease: A Report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. *Circulation.* 2023

AHA/ACC 2021 guidelines on Chest Pain

2021 AHA/ACC/ASE/CHEST/SAEM/SCCT/SCMR Guideline for the Evaluation and Diagnosis of Chest Pain. *Circulation.* 2021

ACC/AHA/SCAI 2021 guideline on revascularization

2021 ACC/AHA/SCAI Guideline for Coronary Artery Revascularization: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines

ESC/EACTS 2018 guidelines on revascularization

2018 ESC/EACTS Guidelines on myocardial revascularization. *Eur Heart J.* 2018.

ESC 2023 Guidelines in diabetes

2023 ESC Guidelines for the management of cardiovascular disease in patients with diabetes. *Eur Heart J.* 2023.

ESC 2016 guidelines on AF.

2016 ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur Heart J* 2016;37:2893-2962.

ACC/AHA/ACCP/HRS 2023 guideline on AF

2023 ACC/AHA/ACCP/HRS Guideline for the Diagnosis and Management of Atrial Fibrillation: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation.* 2023

ACC/AHA 2019 Guideline on the Primary Prevention of Cardiovascular Disease

2019 ACC/AHA Guideline on the Primary Prevention of Cardiovascular Disease. *J Am Coll Cardiol.* 2019;74:1376-1414.

ACC/AHA 2026 Guideline on Management of Dyslipidemia

ACC/AHA/AACVPR/ABC/ACPM/ADA/AGS/APhA/ASPC/NLA/PCNA Guideline on the Management of Dyslipidemia: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation.* 2026.

ESC 2019 Guidelines on Dyslipidaemias

2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J.* 2019.

ESC 2025 Update of Guidelines on Dyslipidaemias

2025 Focused Update of the 2019 ESC/EAS Guidelines for the management of dyslipidaemias. *Eur Heart J.* 2025

ESC 2021 Guidelines on cardiovascular disease prevention

2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J.* 2021

ESC 2022 Guidelines on noncardiac surgery

2022 ESC Guidelines on cardiovascular assessment and management of patients undergoing non-cardiac surgery. *Eur Heart J.* 2022

AHA/ACC 2024 Guideline on noncardiac surgery

2024 AHA/ACC/ACS/ASNC/HRS/SCA/SCCT/SCMR/SVM Guideline for Perioperative Cardiovascular

Management for Noncardiac Surgery. JACC 2024

ESC 2020 Guidelines on sports

Pelliccia A, Sharma S, Gati S, et al. 2020 ESC Guidelines on sports cardiology and exercise in patients with cardiovascular disease. *Eur Heart J*. 2020.

ESC 2022 Guidelines on cardio-oncology

2022 ESC Guidelines on cardio-oncology developed in collaboration with the European Hematology Association (EHA), the European Society for Therapeutic Radiology and Oncology (ESTRO) and the International Cardio-Oncology Society (IC-OS). *Eur Heart J*. 2022.

ESC 2018 Guidelines on pregnancy.

ESC Guidelines on the management of cardiovascular diseases during pregnancy. *Eur Heart J*. 2018.

Relevant Scientific Statements

AHA 2022 Scientific Statement on CVD Risk Factors in Diabetes.

Comprehensive Management of Cardiovascular Risk Factors for Adults With Type 2 Diabetes. A Scientific Statement From the American Heart Association. *Circulation*. 2022 Jan 10;CIR0000000000001040

ACC 2020 Expert Consensus on Diabetes

2020 Expert Consensus Decision Pathway on Novel Therapies for Cardiovascular Risk Reduction in Patients With Type 2 Diabetes: A Report of the American College of Cardiology Solution Set Oversight Committee. JACC 2020; 76: 1117-1145

AHA 2021 statement on obesity and CVD

Obesity and Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation*. 2021

AHA 2021 Statement on Practices in the Cardiac Catheterization Laboratory

Evidence-Based Practices in the Cardiac Catheterization Laboratory: A Scientific Statement From the American Heart Association. *Circulation*. 2021;CIR0000000000000996.

ESC/EACTS review of the 2018 guideline on revascularization

2022 Joint ESC/EACTS review of the 2018 guideline recommendations on the revascularization of left main coronary artery disease in patients at low surgical risk and anatomy suitable for PCI or CABG. *Eur Heart J*. 2023;ehad476.

AHA 2025 Scientific Statement on Alcohol and CVD

Alcohol Use and Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation*. 2025.

AHA 2025 Scientific Statement on Intraoperative Transesophageal Echocardiography During Adult Cardiac Surgery. Considerations of Intraoperative Transesophageal Echocardiography During Adult Cardiac Surgery: A Scientific Statement From the American Heart Association. *Circulation* 2025

AHA 2025 Scientific Statement on “Food is Medicine”

A Systematic Review of "Food Is Medicine" Randomized Controlled Trials for Noncommunicable Disease in the United States: A Scientific Statement From the American Heart Association. *Circulation*. 2025.

AHA 2025 Scientific Statement on Coronary Artery Calcium

Opportunistic Detection of Coronary Artery Calcium on Noncardiac Chest Computed Tomography: An Emerging Tool for Cardiovascular Disease Prevention: A Scientific Statement From the American Heart Association. *Circulation*. 2025;152:e391-e401.

AHA 2025 Scientific Statement on Cardiac Rehabilitation in Women

Cardiac Rehabilitation in Women: A Scientific Statement From the American Heart Association. *Circulation*. 2025;152:e376-e390.

AHA 2026 Scientific Statement on Dietary Guidance

2026 Dietary Guidance to Improve Cardiovascular Health: A Scientific Statement From the American Heart Association. *Circulation*. 2026.

AHA 2026 Scientific Statement on Secondary Prevention After CABG

Secondary Prevention After Coronary Artery Bypass Graft Surgery: 2026 Update: A Scientific Statement From the American Heart Association. *Circulation*. 2026

ESC and EACTS 2026 antithrombotic therapy after CABG

Antithrombotic therapy after coronary artery bypass graft surgery: a Clinical Consensus Statement of the ESC Working Group on Cardiovascular Surgery, the ESC Working Group on Cardiovascular Pharmacotherapy, and the European Association for Cardio-Thoracic Surgery (EACTS). *Eur Heart J*. 2026;47:2419-2436.

ESC groups 2026 Statement on sex differences in IHD.

Sex and gender differences in coronary pathophysiology and ischaemic heart disease. *Eur Heart J*. 2026;47:2727-2749.

ACC 2026 Report on Antiplatelet Therapy

Antiplatelet Therapy in the Management of Atherosclerotic Cardiovascular Disease: 2026 ACC Scientific Statement: A Report of the American College of Cardiology. *J Am Coll Cardiol*. 2026.

AHA 2026 Statement on Caffeine

Caffeine and Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation*. 2026;154:e345-e355.

EPIDEMIOLOGY AND PATHOPHYSIOLOGY OF CORONARY ARTERY DISEASE

Definitions and Classification

Persons with atherosclerosis of epicardial coronary arteries and/or microcirculation may be asymptomatic or present with angina pectoris on effort or develop an acute coronary syndrome (ACS). **Stable coronary artery disease** may be detected following a diagnostic ischaemia test, or get diagnosed after presentation with an acute coronary syndrome. **Acute Coronary Syndrome (ACS)** refers to an acute imbalance of myocardial oxygen supply and demand due to progressive or abrupt flow-limiting coronary stenosis, and/or high-output or increased afterload states. ACS include myocardial infarction with ST segment elevation or new LBBB (**STEMI**), and non-ST-elevation myocardial infarction (**NSTEMI**) that is diagnosed by enzyme rise. **Unstable angina** is diagnosed when there are new or worsening symptoms of ischaemia, and ischemic ECG changes such as ST-segment depressions and T-wave inversion, with normal biomarkers. It is now evident that a large majority of patients with clinical manifestations of myocardial ischaemia, with rest pain but without elevated troponins by a commercially available assay, and therefore considered to have UA, have an elevation of circulating troponins measured by a high-sensitivity assay and could therefore be classified as NSTEMI.¹

Epidemiology

The global burden of Cardiovascular Disease (CVD) is increasing, primarily because of the ageing of the population.² Atherosclerotic cardiovascular diseases were estimated to be responsible for 21% of all deaths in the third *Lancet* Commission on Investing in Health, Global Health 2050 that estimated crude mortality rates from 1970 to 2019.³ Silent atherosclerosis is detectable in early adulthood (18 years of age).⁴

According to NHANES 2011 to 2013 data, the prevalence of CVD, comprising coronary artery disease (CAD), heart failure (HF), stroke, and hypertension) in adults ≥ 20 years of age was 48.9% overall (130.6 million in 2012) and increases with age in both males and females.⁵

The total CHD prevalence is 5.2% in US adults ≥ 20 years of age, affecting 15.9 million Americans. CHD prevalence is higher in males (6.9%) than females (3.7%).⁵ Epidemiology data are presented in Figures 1 and 2. It is estimated that in the USA by the year 2060, compared with the year 2025, the number of people with ischemic heart disease will increase by 31.1% (21.9M to 28.7M), heart failure by 33.0% (9.7M to 12.9M), myocardial infarction

by 30.1% (12.3M to 16.0M), and stroke by 34.3% (10.8M to 14.5M).⁶ CVD remains the most common cause of death in ESC member countries with over 3 million deaths per year.⁷ Although the risk-factor burden is lowest in low-income countries, the rates of major cardiovascular disease and death are substantially higher in low-income countries than in high-income countries, probably due to inferior control of risk factors and proper therapy.⁸ In the developed world, CVD is the single most common cause of death (WHO Report 2020). According to the Global Burden of Diseases, Injuries, and Risk Factors Study (GBD), in 2019, CVD was the underlying cause of 9.6 million deaths among men and 8.9 million deaths among women, approximately one-third of all deaths globally. Of these deaths, 6.1 million occurred in people between the ages of 30 to 70 years. Despite a 30-year decline in age-standardized CVD mortality rates globally, the GBD 2019 study reveals increasing age-standardized CVD mortality in many locations since 2010 and an overall flattening of the global decline in the past 5 years.⁹

Approximately half of CVD mortality is due to CAD.^{10, 11} Atherosclerosis in general can be detected by computed tomography angiography in up to 42% of apparently healthy individuals aged 50-64 years, although significant stenosis ($\geq 50\%$) is less common (5%).¹² Worldwide, more than 3 million people each year are estimated to have a STEMI and more than 4 million have a NSTEMI.¹³ In the USA, the estimated annual incidence of MI is 605 000 new attacks and 200 000 recurrent attacks.⁵ Hospital mortality is higher in patients with STEMI but in the long-term mortality is higher in patients with NSTEMI.¹⁴ MI that occurs in the community still carries a 25% mortality risk, whereas in-hospital mortality without fibrinolysis approaches 15%. Current pharmacological and mechanical therapeutic approaches have reduced this risk to 3-5%.^{13, 15} In-hospital occurring STEMI carries a higher mortality, especially in patients admitted for noncardiac reasons.¹⁶ Silent myocardial infarctions represent >45% of incident myocardial infarctions and are associated with poor prognosis.¹⁷

The estimated direct and indirect cost of IHD in 2017 to 2018 (average annual) was \$228.7 billion. MI (\$12.1 billion) and CAD (\$9.0 billion) were 2 of the 10 most expensive conditions treated in US hospitals in 2013.¹⁸ A study examining public and private health insurer records in the USA from 1996 to 2016 reported an estimated cost of \$28.4 billion for IHD out of a total of 3.1 trillion dollars for health care in general.¹⁹ Between 2013 and 2030, medical costs of CAD are projected to increase by approximately 100%.²⁰ CVD accounts for 11% of the European Union's total healthcare expenditure.⁷

Aetiology

Coronary artery disease is due to a complex interaction of political, social, behavioural, physical, biological, and genetic risk factors.²¹ Noncommunicable diseases such as air and

noise pollution, chemical contamination, artificial light at night, and climate change effects like heat extremes, floods, wildfires, and desert storms, are the leading cause of mortality globally, accounting for 70% of deaths, with cardiovascular disease constituting 44% of these fatalities.²² More than 90% of myocardial infarctions are attributable to modifiable risk factors such as ***hypertension, dietary risks, dyslipidaemia, smoking, diabetes, exposure to traffic air pollution and traffic noise, obesity*** (not in the very elderly) and, especially, ***abdominal obesity, insomnia, psychosocial factors, and very low socioeconomic status***.^{9, 23-32} ***Hypertension*** is the most important modifiable factor (Table 1),³¹ and CV risk increases at a lower BP level in females than in males.³³ ***Smoking*** is associated with a substantially increased risk of myocardial infarction that is reduced but not becoming negligible with electronic cigarettes.³⁴ Noncigarette tobacco use (cigar, pipe, and smokeless tobacco) is associated with increased inflammatory markers related to cardiovascular harm, including high-sensitivity C-RP fibrinogen, and measures of carotid vascular disease.³⁵ ***Dietary*** habits are important. ***Saturated fat and trans-fats*** (mainly hydrogenated oils), high ***carbohydrate intake, processed meat and ultra-processed foods*** have been associated with increased CVD risk and total mortality. ***Ultraprocessed*** foods and beverages in particular, pose a growing public health challenge.^{36, 37} A simplified guide to Nova food classification for clinical use is presented in Table 2. Excess ***sodium*** intake is associated with an increased risk of coronary heart disease and cardiovascular mortality.^{38, 39} The data on ***alcohol*** suggest possible risk reduction when alcohol, especially wine, is consumed in low amounts (such as no more than 1 to 2 drinks a day) in regard to coronary artery disease, stroke, sudden death, and possibly heart failure. ***Coffee*** consumption, 2-3 cups a day is most probably associated with a reduced risk of CVD, whereas high doses of caffeine such as that found in energy drinks may be associated with cardiovascular harm.^{40, 41} For a detailed discussion on alcohol and caffeine see the Chapter on Prevention.

However, the limitations of non-randomized, observational trials on human nutrition are well established,⁴² and focusing on a single lipid marker such as LDL cholesterol alone, does not capture the net clinical effects of nutrients on cardiovascular risk. The evidence from observational studies examining the relationship between dietary ***cholesterol*** and CVD risk is inconsistent, but the discrepant results are likely heavily contributed to by residual confounding, since it is difficult to distinguish between the effect of dietary cholesterol per se and the effect of dietary patterns high in cholesterol or ***saturated fat***.⁴³ In general, studies on dietary habits (“Food is Medicine”) often positively influence diet quality and food security, but the impact on clinical outcomes has so far been inconsistent and often failed to reach statistical significance.⁴⁴

Coronary artery disease has a multifactorial aetiology, causality on certain occasions is a matter of judgement, and risk factors should be considered within the context of

socioeconomic and political factors. In approximately 15% of patients with CAD, no traditional risk factor can be identified.⁴⁵

Lipids

Plasma lipid levels are, to a very large extent, determined by genetic factors. In its more extreme forms this is manifested as familial dyslipidaemias. Patients with extreme hypercholesterolemia have an elevated risk of ischemic events. The **severe hypercholesterolemia phenotype** includes all patients with low-density lipoprotein cholesterol levels >190 mg/dl, regardless of cause. There is now unequivocal evidence for a causal effect of LDL-C.⁴⁶ Incident CVD event risk depends on cumulative prior exposure to LDL-C. As such, individuals with the same cumulative LDL-C exposure at age 40 years but with a greater fraction of that exposure occurring earlier in life (between 18 and 30 years) have a greater risk of subclinical plaque development and incident CVD event risk after age 40 years.⁴⁷ Non-HDL cholesterol concentrations in blood are strongly associated with long-term risk of atherosclerotic cardiovascular disease.⁴⁸ Autosomal dominant hypercholesterolemia results from mutations in genes controlling LDL levels (Table 3). This includes familial hypercholesterolemia, as well as other inherited forms of a potential polygenic origin.⁴⁹ Rare variations in a gene (APOC3) that encodes for apolipoprotein C3 are associated with reduced triglyceride levels and lower risk for heart disease.^{50, 51} There is a causal effect of **triglycerides** on CAD risk, but the role for **HDL-C** though possible, remains less certain, particularly as far as therapeutic interventions are concerned.⁵²⁻⁵⁶ In a Danish registry, the association between **HDL cholesterol** concentrations and all-cause mortality was U-shaped, and men and women from the general population with extreme high HDL cholesterol had high all-cause mortality.⁵⁷ However, low-density lipoprotein cholesterol is associated with near-term atherosclerotic cardiovascular disease events in middle-aged patients with presence of coronary atherosclerosis, whereas no association is established for those without coronary atherosclerosis.⁵⁸ **Lipoprotein A Lp(a)**, especially >50 mg/dl (105 mmol/L), is associated with cardiovascular risk continuously and independently of LDL-C level.⁵⁹⁻⁶⁴ Concentrations are more than 90% genetically determined and 17% higher in post-menopausal women than in men.⁶⁵ Blood levels of **apolipoprotein B** (apoB) provide additional atherosclerotic risk information beyond LDL-C across the entire range of LDL-C level.⁶⁶ ApoB measurement in young adulthood provides a comprehensive measurement since it also captures low-density lipoprotein particle (LDL-P), and triglyceride-rich lipoprotein particle (TRL-P) that are associated with CVD risk.^{67, 68} **Remnant cholesterol**, ie total cholesterol minus LDL and HDL, and including the cholesterol content of the triglyceride-rich very-low-density lipoproteins, intermediate-density lipoproteins, and chylomicron remnants in the nonfasting state, is an important risk factor for future cardiovascular events.⁶⁹

Familial hypercholesterolemia (FH) is an autosomal codominant monogenic disorder, characterized by elevated low-density lipoprotein cholesterol (LDL-C) mainly caused by loss-of function mutations in genes related to reduced clearance of LDL particles. It is very rarely recessive, and the phenotype may also be polygenic.⁷⁰⁻⁷² Heterozygous FH (HeFH) affects around 1 in 300 people worldwide, and homozygous FH (HoFH) affects around 1 in 300 000. The prevalence of heterozygous FH (HeFH) is 1:300 in the general population, with data suggesting that $\approx 1\%$ of those potentially affected are detected.⁷³ Homozygous FH (HoFH) affects around 1 in 300 000.⁷⁴ Monogenic familial hypercholesterolemia is associated with lower responsiveness to conventional low-density lipoprotein cholesterol lowering therapies and increased preclinical coronary atherosclerosis when compared with hypercholesterolemia of polygenic or unknown genetic cause.⁷⁵

Familial hypercholesterolemia is clinically diagnosed by five major criteria including: presence of early CAD in the index case, elevated LDL-C ($>190\text{mg/dl}$ - 5 mmol/l), family history of premature CAD (women <60 years, men <55 years), tendon xanthomas or corneal arcus (age <45 years).⁷⁶ The 2015 EAS consensus statement on FH in children proposed that any child with an LDL-C $\geq 5.0\text{ mmol/L}$ (190 mg/dL) measured twice after dietary intervention has a high probability of genetically based FH; this threshold was lowered to $\geq 4.0\text{ mmol/L}$ (155 mg/dL) if there was a family history of premature ASCVD and/or high baseline cholesterol in one parent, and further lowered to LDL-C $\geq 3.5\text{ mmol/L}$ (135 mg/dL) if one parent had a genetic diagnosis of FH.⁷⁷ However, recent data digest that when using these criteria, early diagnosis in almost half of children with an FH-causing variant would have been missed, and a 1016 a European Atherosclerosis Society consensus statement proposed a modification of these criteria.⁷⁴ From 1 year of age, LDL levels discriminate between familial and non-familial hypercholesterolemia in children.⁷⁸

Heterozygous FH is caused by a variant in a single allele of LDLR (LDL receptor), apoB (apolipoprotein B), or PCSK9 (proprotein convertase subtilisin/kexin type 9), whereas **homozygous FH** is produced by variant in 2 alleles of the same or different canonical genes.⁷⁹ Patients with heterozygous FH have a total cholesterol $350\text{--}550\text{ mg/dL}$ ($9\text{--}14\text{ mmol/L}$) and LDL-C values $200\text{--}400\text{ mg/dL}$ ($5\text{--}10\text{ mmol/L}$) (Table 2).^{80 76} In homozygous FH, plasma LDL-C levels are $>10\text{ mmol/L}$ (400 mg/dL).⁸¹ The consequent diminution in LDL receptor activity leads to extreme elevations of LDL cholesterol from birth, accelerated atherosclerosis, and premature death (18 years of age or less) from acute coronary syndromes, often associated with valvar and supra-aortic atheroma of the aortic root. The prevalence of molecularly defined homozygous autosomal dominant familial hypercholesterolaemia is rather high ($1/300,000$), and the clinical phenotype is more variable than previously assumed, with disease often underdiagnosed between familial and non-familial hypercholesterolaemia in children.⁸² **Familial combined hyperlipidaemia** is a highly prevalent mixed

dyslipidaemia (1:100-200) characterized by elevated levels of LDL-C, TGs, or both, and is an important cause of premature CAD.⁸³ In normolipidaemic patients, post-prandial TG values rarely exceed 400 mg/dL (4.6 mmol/L).⁵³ Extremely high fasting triglyceride levels > 880 mg/dL indicate persistent chylomicronemia that is associated with the risk of recurrent acute pancreatitis.⁸⁴ Criteria have been proposed for diagnosis (Table 3) that is confirmed by genetic testing.

Genetic testing is recommended for patients with definite or probable FH, as well as for their at-risk relatives (Table 4).⁸¹ Testing should mainly include the genes encoding the low-density lipoprotein receptor (*LDLR*), apolipoprotein B (*APOB*), and proprotein convertase subtilisin/kexin 9 (*PCSK9*).^{72, 85} Among participants with LDL cholesterol ≥ 190 mg/dL, in general, gene sequencing identifies an FH mutation in <2%. However, in patients presenting with an ACS and LDL-C levels ≥ 160 mg/dL, the prevalence of genetically confirmed FH is high (approximately 9%).^{86, 87} Genetic (polygenic) risk scores also appear valuable in predicting recurrent cardiovascular events in patients, and thus identifying patients in whom PCSK9 therapy is beneficial.^{88, 89}

Comorbidities

Arterial hypertension is the top modifiable risk factor for CVD burden, followed by **dyslipidaemia (high LDL)**.^{9, 31} **Type 2 diabetes mellitus** is an established risk factor for CVD. CVD is the leading cause of death in adults with diabetes mellitus, and the life expectancy of these patients, especially those with a previous MI, is considerably shorter than non-diabetics.⁹⁰ **Chronic kidney disease** is one of the strongest risk factors for the development of CVD. Chronic kidney disease mimics an accelerated aging of the cardiovascular system by causing a systemic, chronic proinflammatory state contributing to vascular and myocardial remodeling processes resulting in atherosclerotic lesions, vascular calcification, and vascular senescence as well as myocardial fibrosis and calcification of cardiac valves.^{91, 92} **Polycystic ovarian syndrome** is associated with increased cardiovascular morbidity and a higher risk of atherosclerotic cardiovascular disease.⁹³ **Obesity** leads to the development of cardiovascular disease independently of other cardiovascular risk factors. However, susceptibility to obesity-related cardiovascular complications is not mediated solely by overall body fat mass and body mass index (BMI), but depends largely on individual differences in regional body fat distribution, such as abdominal obesity and visceral adiposity.^{94, 95} In the elderly (≥ 80 years) both higher BMI and waist circumference are associated with a lower risk of all-cause mortality, but persons with higher BMIs but lower waist circumferences have the lowest mortality risk.⁹⁶ **Congenital heart disease** is an independent risk factor for the occurrence and outcome of myocardial infarction.⁹⁷ A **myocardial bridge**, ie muscle overlying the intramyocardial segment of the epicardial coronary artery, usually the LAD, is a usually benign abnormality of multifactorial origin that, however, may lead to ischaemia.^{98, 99} The

reported prevalence varies widely depending on the imaging modality, appearing in 5%–8% of coronary angiograms and 18%–25% of CCTAs. Autopsy studies, however, suggest an even higher prevalence of 30%–86%, indicating that MB may be more of an anatomical variant than a true anomaly.⁹⁹ **Hyperuricaemia** may contribute to the development and progression of chronic kidney disease (CKD), and, probably, cerebral and cardiovascular disease, and mortality. However, the clinical utility of uric acid-lowering for the prevention or treatment of hypertension, CVD, or CKD is controversial.¹⁰⁰ **Non-alcoholic fatty liver disease** increases the risk of CVD due to systemic inflammation, endothelial dysfunction, hepatic insulin resistance, oxidative stress, and altered lipid metabolism.¹⁰¹ Clinically, both **hyper-** and **hypothyroidism** can affect dyslipidaemia and atherosclerotic vascular disease.¹⁰² Patients with **cancer** have a substantially increased short-term risk of arterial thromboembolism.¹⁰³ **Chronic venous insufficiency** is related to cardiovascular disease and is predictive of all-cause death.¹⁰⁴ Inflammatory and immune mechanisms are implicated in atherosclerosis.¹⁰⁵ **Chronic inflammation** is associated with atherosclerosis as expressed by traditional markers such as the neutrophil-lymphocyte ratio (>2.5), CRP, IL-1 and IL-6.^{62, 106} A hrsCRP >3 mg/L indicates an increased cardiovascular risk in women, even in the absence of traditional risk factors such as hypertension, dyslipidemia, diabetes and smoking.¹⁰⁷ **Severe infections** that require hospitalization are associated with increased risks for major cardiovascular disease events.¹⁰⁸ Chronic inflammatory diseases including **HIV** infection, **psoriasis**, **rheumatoid arthritis**, and **systemic lupus erythematosus** predispose to atherosclerotic cardiovascular disease.^{109–111} **Hodgkin lymphoma** survivors have an increased risk of cardiovascular disease for at least 40 years after diagnosis.¹¹² Patients with **antiphospholipid syndrome** are at increased risk for accelerated atherosclerosis, myocardial infarction, stroke, and valvular heart disease.¹¹³ However, factor V Leiden is not associated with increased risk of atherothrombotic events and mortality in patients with established and treated CAD.¹¹⁴ **Kawasaki disease** is an acute vasculitis of childhood that leads to coronary artery aneurysms in ≈25% of untreated cases.¹¹⁵ **Community-acquired bacteremia** is associated with increased short-term risk of myocardial infarction and stroke.¹¹⁶ **Respiratory infections**, especially influenza, are associated with acute myocardial infarction.¹¹⁷ Patients with **syphilis** have a higher risk of cardiovascular events and all-cause mortality.¹¹⁸ **COVID-19 pneumonia** is associated with acute myocardial injury and thrombosis.^{119, 120} **Herpes zoster** infection is associated with CVD.¹²¹ **Fibromuscular dysplasia** involving the coronary arteries is an uncommon but important condition that can present as acute coronary syndrome.^{122, 123} **Endometriosis** is associated with a higher risk of cardiovascular events including myocardial infarction, arrhythmia, and heart failure.¹²⁴ **Periodontitis** is more common in patients with a myocardial infarction, and it seems that the relation periodontal disease and CVD is causal.¹²⁵ The prevalence of **obstructive sleep apnoea** is as high as 40% to 80% in patients with

hypertension, heart failure, coronary artery disease, pulmonary hypertension, atrial fibrillation, and stroke.¹²⁶ **Mental and substance addiction disorders**, especially when diagnosed at a young age, are associated with increased risk of cardiovascular disease.¹²⁷

Pregnancy complications such as spontaneous preterm delivery, miscarriages, placenta abruption and still birth are associated with increased rates of future CVD, cerebral infarction, and renovascular hypertension.^{128, 129} This has been attributed to either shared etiology or the initiation of pathological processes by a pregnancy loss leading to atherosclerosis.

Drugs

Chemotherapeutic agents for cancer may also cause accelerated atherosclerosis and arterial thrombotic events (see Tables on Risk Assessment). For a detailed discussion see Chapter 92. **NSAIDs** (both non-selective and selective COX-2 inhibitors that spare COX-1 in the gut) are associated with an increased thrombotic risk (see Chapter 30).¹³⁰ Platelet cyclo-oxygenase (COX)-1 activity results in production of thromboxane A, whereas COX-2 activity results in the production of prostacyclin (PGI₂). Regular acetaminophen has been associated with hypertension,¹³¹ and **sodium-containing acetaminophen** (e.g. the effervescent or soluble) is associated with increased risks of CVD and all-cause mortality among individuals with or without hypertension.¹³² Women who use **oral contraceptives containing ethinyl estradiol** are at dose-dependent increased risk for thrombotic stroke and myocardial infarction compared to women using progesterone only contraception.¹³³ These cardiovascular risk factors induce oxidative stress that causes endothelial inflammation and damage. Both **androgen deprivation therapy**, prescribed for prostate cancer, and **testosterone therapy** has been associated with increased risk of CVD.^{134, 135 130} Long-term **anabolic-androgenic steroids** use is associated with increased mortality due to cardiovascular disease.^{136, 137} **Antipsychotic drugs** use may also be associated with a transient increase in risk for AMI, possibly mediated by dopamine type 3 receptor blockades.¹³⁸ **Clarithromycin** has been found to increase cardiovascular mortality in patients with stable coronary heart disease who were not on statin,¹³⁹ and in February 2018 the FDA issued a safety communication. However, the evidence is rather inconsistent. **Opioid** use has been associated with acute coronary syndromes.¹⁴⁰ **Cocaine** has a multitude of pathophysiological pathways by which it affects the cardiovascular system, including coronary vasoconstriction and acute myocardial infarction.¹⁴¹

Other conditions

Environmental risk factors are emerging as critical contributors to noncommunicable diseases.²² **Traffic air pollution** (mainly fine particles especially <2.5 µm in diameter-PM_{2.5}, as well as PM₁₀, CO, NO₂, and ozone) as well as **microplastics** (defined as particles smaller than 5 mm) and **nanoplastics** (particles smaller than 1000 nm) that are released

by plastics degradation are associated with cardiovascular disease.¹⁴²⁻¹⁴⁵ **Radiation-associated cardiac disease** is a heterogeneous and complex disease, with cardiac manifestations which range from pericarditis, coronary artery disease, valvular heart disease, nonischemic cardiomyopathy, and conduction abnormalities, and may occur years or even decades following radiation exposure to the chest.¹⁴⁶ Both **early** and **late menarche** are associated with increased risk of CAD.¹⁴⁷ Self-reported symptoms of **anxiety** and **depression** as well as outbursts of **anger and emotional upset** increase the risk of myocardial infarction.^{148 149 150} **Psychological stress** is a risk factor and prognostic factor for CAD,¹⁵¹ and psychosocial factors in youth such as the socioeconomic and emotional environments predict development of CAD.¹⁵² **Short sleep duration** and deprivation (<6h a day) are associated with increased risk of CVD.^{153 154} **Combat experience** is a risk factor for CAD.¹⁵⁵ Cardiovascular disease is a major cause of death among **homeless** adults.²⁸ **Extreme hot and cold temperatures** are associated with increased risk of death from any cardiovascular cause, ischemic heart disease, stroke, and heart failure.¹⁵⁶ **Shorter height** is associated with increased risk for coronary artery disease.¹⁵⁷ The rate of progression of cellular **ageing** in late midlife (reflected by the rate of leucocyte telomere length attrition) relates to vascular damage, independently from contribution of CV risk factor exposure.¹⁵⁸ The possibility of **deep space radiation** promoting endothelial dysfunction and atherosclerosis has also been raised.¹⁵⁹ Non-linear Mendelian randomization analyses suggest an association of 25(OH)D with CVD risk, with increased CVD risk largely restricted to individuals with **low vitamin D** status.¹⁶⁰

Genetic factors

Coronary artery disease is the most studied cardiovascular phenotype for polygenic risk scores.¹⁶¹ However, apart from genetic disorders of lipoprotein metabolism, distinct morphological characteristics such as left main or proximal disease also display a high heritability.¹⁶² Epidemiological studies have shown that genetic predisposition accounts for 40-60% of the risk for coronary artery disease. Genome-wide association studies have led to the discovery of at least 67 loci associated with the risk of coronary artery disease, explaining approximately 15% of the disease's heritability.¹⁶³ Approximately 10 of them are associated with hypertension or hyperlipidemia, and the rest of them mediate their risks through unknown mechanisms. The strongest genetic effect on the risk of coronary artery disease and myocardial infarction is that of single nucleotide polymorphisms at **chromosome 9p21.3** (CDKN2A/2B), that is also a significant risk factor for abdominal aortic aneurysm, stroke, and Alzheimer's disease.¹⁶⁴⁻¹⁶⁶ The 9p21.3 risk allele is carried by 75% of the European population and confers risk of coronary atherosclerosis in a dose-dependent way (by means of allele copies). However, in patients with established CAD, variants at the 9p21 locus do not affect risk of recurrent or subsequent events, including mortality, ie cannot be used for risk stratification purposes.¹⁶⁷ The identification of the **ABO blood group locus** as a risk factor for

myocardial infarction might provide an explanation on why blood group O offers protection from myocardial infarction and thromboembolism compared to blood groups A and B.¹⁶⁸ Common genetic influences appear to affect both height and CAD risk, partly mediated by genetic effects on LDL cholesterol and triglycerides. Genetically predicted short sleep duration is associated with increased risk of CVD.¹⁵³ The presence of a common **Y chromosome variant**, haplogroup I, has been found to increase coronary risk by 50%.¹⁶⁹ This might explain the higher prevalence of CAD in men. All these genetic variants are not yet ready for routine testing. Loss of function mutations in **ANGPTL4**, a protein that inhibits lipoprotein lipase-LPL and thus reduce triglycerides levels and elevate HDL cholesterol, confer lower risk of coronary artery disease.^{170, 171} Loss of function variants of **ANGPTL3**, a LPL inhibitor that reduces triglycerides levels as well as LDL and HDL cholesterol, or treatment antisense oligonucleotides that targeted ANGPTL3 messenger RNA has also retarded the progression of atherosclerosis in mice.^{172, 173} Genetic scores, such as the metaGRS, derived from large biobanks have been developed as predictors of CAD.¹⁷⁴ **TSR1**, a novel gene located in the chromosome 17p13.3 region, has been identified as a potential causal gene of spontaneous coronary artery dissection.¹⁷⁵ **JCAD** (Junctional cadherin 5 associated, also known as *KIAA1462*), also encodes a junctional protein associated with CAD.¹⁷⁶ A locus on chromosome 1p21.3 harbouring choline-like transporter 3 (**SLC44A3**), has been implicated in the pathophysiology of vulnerable plaques and development of MI.¹⁷⁷ Finally, genetic variants known to affect educational attainment may have broad implications for a health-conscious lifestyle later in life and the risk of CAD.¹⁷⁸ The development of large disease-based consortia and biobanks have enabled the discovery of genetic variants associated with CAD. The availability of these large datasets, along with computational advances, has allowed for the easier derivation, calculation, and validation of genetic (polygenic) risk scores that predict recurrent cardiovascular events in patients with pre-existing CAD.¹⁷⁹

Pathophysiology

Originally thought to be dominantly a lipid storage disease, our current understanding of the pathogenesis of atherosclerosis implicates endothelial injury and inflammation. Following endothelial injury, smooth muscle cells and monocytes/macrophages migrate into arterial intima in the preatherosclerotic stage and during plaque development, and become foam cells by accumulating cholesterol.¹⁸⁰ Additional chemical and mechanical factors that may trigger endothelial injury are altered sheer stress, high oxidative stress, smoking, and insulin resistance.¹⁸¹ More than half of men and women with suspected CAD have microvascular dysfunction even with the absence of an abnormal stress test, and this is a major predictor of

major adverse events.¹⁸² Coronary microvascular disease may represent an epiphenomenon or contribute to the pathogenesis of cardiovascular disease. The development of atherosclerotic disease may be an age-related asymptomatic process or lead to stable ischemic heart disease or an acute coronary syndrome.

Stable ischemic heart disease

Non-acute myocardial ischaemia syndromes are caused by ischaemia due to epicardial coronary stenosis or by ischaemia with non-obstructive coronary arteries (**INOCA**).¹⁸³ The underlying mechanism of chronic CAD is the chronic limited ability to increase oxygen supply to the myocardium in the setting of increased oxygen demand, usually caused by the obstruction of at least one large epicardial coronary artery by atheromatous plaque, microvascular dysfunction, vasoconstriction at the site of dynamic stenosis, or combination of the above. Because myocytes already extract about 75% of the oxygen in coronary blood at rest, a higher demand is primarily met by increasing coronary blood flow. Myocardial ischaemia results in hypoxia that activates cellular anaerobic pathways; produced mediators such as lactate are responsible for the sensation of pain. The development of **collateral circulation** is an adaptive response to ischemia, and may be partially responsible for the nonlinear relationship between stenosis severity, ischemia, and symptom burden.¹⁸⁴

Up to 70% of patients presenting to the cardiac catheterization laboratory with angina may have no angiographic evidence of coronary artery disease (**Ischaemia with nonobstructive coronary arteries – INOCA**), and an increasing number of patients, especially women, develop signs and symptoms of myocardial ischaemia in the absence of obstructive coronary artery disease.^{182, 185-187} In the CORONER study on stable CAD outpatients, incident MI occurred at a stable rate of 0.8% annually, and was related to late stent thrombosis in one-fifth of cases.¹⁸⁸

Acute coronary syndromes

Acute myocardial ischaemia syndromes are caused by coronary plaque rupture or erosion or by myocardial infarction with non-obstructive coronary arteries (**MINOCA**).¹⁸³ The most frequent mechanism of an ACS is a reduction of the myocardial oxygen supply usually due to rupture or erosion of a vulnerable atherosclerotic plaque, which results in endothelial injury and associated thrombosis and dynamic vasoconstriction. Vulnerable plaques likely to rupture or erode have evidence of inflammation with macrophages that overproduce matrix-metalloproteinases (MMP) with collagenase activity, and T-cell infiltrates.¹⁸⁹ Histologically, vulnerable plaques have thin fibrous caps (<65 µm) with large lipid cores (thin-cap fibroatheroma-TCFA)¹⁹⁰⁻¹⁹³ although statin therapy in the modern era may modify their characteristics.¹⁹⁴ The detection of lipid accumulation within the coronary artery wall predicts cardiovascular outcomes.¹⁹⁵ Usually, atherosclerosis occurs within usually in an environment

of low shear stress, but high shear stress may promote plaque rupture.¹⁹⁶ Plaque rupture remains the leading cause of ACS, accounting for approximately 60%–70% of cases.¹⁹⁷ Plaque rupture in non-obstructive lesions of non-infarct related arteries is present in 12% of patients with acute MI. More than half of untreated ruptures transitioned into stable morphologies.¹⁹⁸ Plaque erosion (underlying between 25% to 45% of ACS), intraplaque hemorrhage, calcified nodule, and functional coronary alterations also play an important triggering role.^{199 197, 200} Plaque haemorrhage may occur from plaque rupture (fissure) or from neovascularization (angiogenesis). Plaque fissure or rupture exposes the highly thrombogenic subendothelium (collagen and tissue factor) to circulating platelets and white blood cells. Platelets are also activated by vasoconstrictors such as thromboxane A₂, and adenosine diphosphate (ADP) that binds to P₂Y₁₂ receptors, and cause increased expression of glycoprotein IIb/IIIa that binds fibrinogen or von Willenbrand factor. Thus, platelet aggregation and creation of the *white clot* occurs. The platelet-derived mediators together with other potent vasoconstrictors such as activated thrombin, oxygen-derived free radicals and endothelin result in dynamic vasoconstriction and transient thrombosis. The coagulation cascade may be activated and factor X leads to generation of thrombin that converts fibrinogen to fibrin (*red clot*).²⁰¹ The term ‘calcified nodule’ indicates calcified coronary atherosclerotic plaque nodules. Fragmentation of the necrotic core of the plaque nodules promotes fibrous cap disruption and luminal thrombosis.²⁰²

It should be noted that erosion or rupture of the coronary atherosclerotic plaque is typically required for an event to happen, but only when coinciding with a prothrombotic state and subsequent thrombosis at the site of plaque rupture or erosion, it leads to an event. In the majority of cases, plaque ruptures or erosions take place in the absence of symptoms and commonly lead to healing and progression of coronary arterial narrowing.²⁰³ Vulnerable plaques, intact or ruptured, may be found along the coronary tree and the true culprit lesion can be difficult to define.²⁰⁴ ACS may also occur without apparent epicardial coronary thrombus or stenosis. Such events may arise from coronary artery spasm, microvascular disease, or other pathways.¹⁸⁹

If the ischaemia is prolonged and severe enough to cause limited myocardial necrosis the diagnosis of **NSTEMI** is put; further damage, usually due to formation of a red thrombus that causes ST elevation, produces the clinical picture of **myocardial infarction (STEMI)**. Thrombosis and occlusion of the coronary artery is the commonest cause of STEMI.^{205 206} Myocardial injury can result from a wide variety of ischemic and nonischemic mechanisms.²⁰⁷ Angiography at the acute phase reveals an occluded coronary artery in 90% of cases, although the converse is not the case due to the possible presence of collaterals. Intracoronary thrombi are mainly composed of fibrin (increased content with increased ischemic time), platelets, erythrocytes, and few cholesterol crystals and leukocytes.²⁰⁵ There is a circadian periodicity

with most events occurring between 6 am and noon. Platelet hyper-reactivity and pro-coagulant states contribute to the thrombotic process that diminishes microcirculatory perfusion by reduced coronary artery flow through epicardial stenoses, as well as by distal embolization of thrombus. Histochemical stains can identify zones of necrosis 2-3 hours after the onset of necrosis, while gross pathology changes can be identified 6-12 hours following occlusion. Eight to 10 days after infarction the thickness of the affected ventricular wall is reduced as necrotic muscle is removed by mononuclear cells, and over the next 2-3 months the firm scar develops. Infarct expansion (disruption and tissue loss not explained by additional necrosis) and ventricular dilation may also occur and determine ventricular remodeling. The systemic inflammatory reaction to acute myocardial infarction can aggravate inflammation in the plaque, thus resulting in recurrent thrombotic events in the aftermath of an acute coronary syndrome.¹⁸¹ Following reperfusion, the course of myocardial edema is bimodal. Acute reperfusion is responsible for the initial acute edematous reaction that dissipates at 24 h, and then initiates days after infarction, reaching a plateau at 4-7 days because of tissue healing processes.^{208, 209} In the modern era of primary PCI, improved LVEF is detected in 55% of patients one month after the event.²¹⁰

Myocardial infarction is now classified into one of three clinical types: primary, secondary and procedure related (see also Chapter on Diagnosis of ACS). **Primary myocardial infarction** is due to spontaneous presentation due to a primary acute coronary pathology. The diagnosis is confirmed and aetiology defined if atherothrombosis or an alternative acute coronary pathology is identified. **Secondary myocardial infarction results** from myocardial oxygen supply–demand imbalance due to another acute condition. However, two-thirds of these patients may have coronary artery disease, and one-third had left ventricular impairment, which were previously unrecognized and untreated in the majority. Thus, investigation with invasive or noninvasive coronary and cardiac imaging should be considered in patients with secondary infarction.²¹¹ **Procedure-related myocardial infarction** occurs as a complication of a percutaneous or surgical cardiac procedure.

Primary myocardial infarctions have been previously shown to frequently occur at sites of mild-to-moderate stenosis.²¹²⁻²¹⁶ Postmortem and older angiographic data, however, have demonstrated that ruptured plaques leading to thrombosis more likely occur within the segment of complex lesions with significant (>50%) stenoses.²¹⁷⁻²²¹ In the PROSPECT trial on patients with acute coronary syndromes treated by coronary intervention, 3-year recurrence rates were 20%. Half of them occurred in angiographically mild (<70% stenosis), non-culprit lesions. However, predictors of recurrent events were both a luminal area ≤ 4 mm², and the presence of thin-cap fibroatheroma.²²² An analysis of the data of the COURAGE trial has also shown that MIs are not more likely to occur from mild coronary lesions compared with moderate or severe ones; the number of non-revascularized lesions

$\geq 50\%$ was the only angiographic predictor of an acute coronary syndrome.²²³ In the ROMICAT trials, lesions with $\geq 50\%$ stenosis were detected in 78% of patients with ACS.²²⁴ Data from a large retrospective cohort study of US veterans undergoing elective coronary angiography, have also demonstrated that obstructive (stenoses $\geq 70\%$) CAD was associated with worse outcomes such as MI and death. Patients with nonobstructive CAD were at increased risk compared to those with no apparent CAD.²²¹ In the SCOT-HEART trial, adverse coronary plaque characteristics and overall calcified plaque burden conferred an increased risk of coronary heart disease death or nonfatal myocardial infarction.²²⁵ In a Danish CT angiography study on asymptomatic persons, subclinical, obstructive ($>50\%$ diameter stenosis) coronary atherosclerosis was associated with a more than 8-fold elevated risk for myocardial infarction.²²⁶

Thin-cap fibroatheroma (TCFA) lesions with a large plaque burden as detected by intravascular ultrasound radiofrequency analysis, carry higher risk than small thin-cap fibroatheroma lesions.²²⁷ When intravascular imaging information is available, the lesions with $<85\mu\text{m}$ thick cap and $>75\%$ cross-sectional luminal area stenosis are the most probable to rupture.¹⁹¹ The absolute number of TCFA has been found 3 times greater in nonsevere stenosis than in severe stenosis. It was, however, twice as likely for a lesion to be TCFA in cases of severe stenosis than in nonsevere stenosis, and TCFA in severely-stenotic areas had more features of plaque vulnerability.²²⁸ Nonculprit lesions identified by optical coherence tomography (OCT), as thin-cap atheroma and/or having a minimum luminal area $<3.5\text{ mm}^2$, many of which appear angiographically mild, were also associated with a high risk of future ischemic events.¹⁹² Plaque healing may occur, and the evolution of coronary artery disease appears to be more stable in patients with healed plaques than in patients with unhealed plaques.²²⁹

Additional causes of UA/NSTEMI are coronary spasm, coronary emboli, coronary arterial inflammation, or spontaneous dissection. Spontaneous **coronary artery dissection** is frequent cause of myocardial infarction (up to 35%) in the young, and especially women <50 years of age. Postpartum status, fibromuscular dysplasia, cocaine use, Ehlers-Danlos syndrome, hormonal therapy, connective tissue disorders, systemic inflammatory disease, and extreme physical exertion are conditions that have been associated with spontaneous coronary dissection.²³⁰⁻²³⁴ There has also been some evidence that SCAD may be caused by intramural bleeding (the outside-in mechanism), possibly associated with more potent antiplatelet therapy.²³⁵ Hypertension increases the risk of recurrent dissection, whereas beta-blocker therapy is protective. Pregnancy-related spontaneous coronary artery dissection is the most common cause of pregnancy-associated myocardial infarction.²³⁶ It is estimated that 1 in 16000 pregnancies is complicated by acute myocardial infarctions in the US, and up to one

fourth of these are due to spontaneous coronary dissection.^{237, 238} It is a rare and potentially lethal complication of pregnancy.

Epicardial or microvascular coronary spasm occurs commonly in the absence of severe coronary disease and can present with or without anginal pain.²³⁹ It may be associated with exogenous or endogenous catecholamines excess.²⁴⁰ Epicardial coronary spasm is a well-reported cause of cardiac arrest. Identification of non-critical coronary disease in the absence of other causes of cardiac arrest might suggest the diagnosis, particularly in cocaine abusers, as well as in smokers.²⁴¹⁻²⁴³

Coronary embolism occurs in up to 3% of patients with STEMI, and AF is the most frequent cause.²⁴⁴

Eosinophilic myocarditis is also a cause of coronary vasculitis and thrombosis.²⁴⁵

Myocardial infarction with nonobstructive coronary arteries (MINOCA) occurs in 5% to 10% of all patients with acute MI and is usually seen in younger patients and women.^{246, 247} The most common conditions associated with MINOCA are epicardial coronary artery disease with possible contribution of coronary artery spasm (e.g. plaque rupture with transient thrombosis), imbalance between oxygen supply and demand (e.g. coronary artery spasm and coronary embolism), microvascular dysfunction, hereditary thrombophilia disorders, and secondary to myocardial disorders without involvement of the coronary arteries (e.g. myocarditis or Takotsubo cardiomyopathy).²⁴⁶⁻²⁴⁸ Most patients clinically diagnosed with MINOCA have a final diagnosis of myocardial infarction (59%), and an alternate diagnosis mimicking myocardial infarction is present in 20%.²⁴⁹ Most patients (up to 64%) have a final diagnosis of myocardial infarction.²⁴⁸ Approximately 20% of patients with MINOCA and age >65 years in a MEDICARE population suffered a major adverse event over the next 12 months.²⁵⁰ In the CIAO group of the ISCHAEMIA study, half of the patients had resolution of stress-induced wall motion abnormalities on repeat stress echocardiography at 1 year and symptoms improve in 43%. There was no significant correlation between change in angina and change in ischaemia on stress echocardiography.²⁵¹

Stunned myocardium refers to depressed contractility following transient ischaemia prior to full recovery, and having sustained no irreversible myocyte damage.²⁵² **Hibernating myocardium** is dysfunctional and viable myocardium, but with more severe cellular structural changes, that recovers systolic function following revascularization.²⁵²

References

1. Braunwald E and Morrow DA. Unstable angina: is it time for a requiem? *Circulation*. 2013;127:2452-7.
2. Vaduganathan M, et al. The Global Burden of Cardiovascular Diseases and Risk: A Compass for Future Health. *J Am Coll Cardiol*. 2022;80:2361-2371.
3. Chang AY, et al. Epidemiological and demographic trends and projections in global health from 1970 to 2050: a descriptive analysis from the third Lancet Commission on Investing in Health, Global Health 2050. *Lancet*. 2025;406:940-949.
4. Bundgaard H and al. e. Prevalence of Silent Atherosclerosis across Adult Life. *NEJM*. 2026.
5. Palaniappan LP, et al. 2026 Heart Disease and Stroke Statistics: A Report of US and Global Data From the American Heart Association. *Circulation*. 2026;153:e275-e906.
6. Mohebi R, et al. Cardiovascular Disease Projections in the United States Based on the 2020 Census Estimates. *J Am Coll Cardiol*. 2022;80:565-578.
7. Timmis A, et al. European Society of Cardiology: the 2023 Atlas of Cardiovascular Disease Statistics. *Eur Heart J*. 2024;45:4019-4062.
8. Yusuf S, et al. Cardiovascular risk and events in 17 low-, middle-, and high-income countries. *N Engl J Med*. 2014;371:818-27.
9. Roth GA, et al. Global Burden of Cardiovascular Diseases and Risk Factors, 1990-2019: Update From the GBD 2019 Study. *J Am Coll Cardiol*. 2020;76:2982-3021.
10. Heart Failure and Cardiovascular Diseases *European Heart Network*. 2019.
11. Timmis Chair Writing Group A, et al. European Society of Cardiology: cardiovascular disease statistics 2021. *Eur Heart J*. 2022;43:716-799.
12. Bergstrom G, et al. Prevalence of Subclinical Coronary Artery Atherosclerosis in the General Population. *Circulation*. 2021;144:916-929.
13. White HD and Chew DP. Acute myocardial infarction. *Lancet*. 2008;372:570-84.
14. Terkelsen CJ, et al. Mortality rates in patients with ST-elevation vs. non-ST-elevation acute myocardial infarction: observations from an unselected cohort. *Eur Heart J*. 2005;26:18-26.
15. Puymirat E, et al. Acute Myocardial Infarction: Changes in Patient Characteristics, Management, and 6-Month Outcomes Over a Period of 20 Years in the FAST-MI Program (French Registry of Acute ST-Elevation or Non-ST-Elevation Myocardial Infarction) 1995 to 2015. *Circulation*. 2017;136:1908-1919.
16. Garberich RF, et al. ST-elevation myocardial infarction diagnosed after hospital admission. *Circulation*. 2014;129:1225-32.
17. Zhang ZM, et al. Race and Sex Differences in the Incidence and Prognostic Significance of Silent Myocardial Infarction in the Atherosclerosis Risk in Communities (ARIC) Study. *Circulation*. 2016;133:2141-8.
18. Tsao CW, et al. Heart Disease and Stroke Statistics-2022 Update: A Report From the American Heart Association. *Circulation*. 2022;145:e153-e639.
19. Dieleman JL, et al. US Health Care Spending by Payer and Health Condition, 1996-2016. *JAMA*. 2020;323:863-884.
20. Coronary Revascularization Writing G, et al. ACC/AATS/AHA/ASE/ASNC/SCAI/SCCT/STS 2016 Appropriate Use Criteria for Coronary Revascularization in Patients With Acute Coronary Syndromes. *J Am Coll Cardiol*. 2016.
21. Gupta R and Wood DA. Primary prevention of ischaemic heart disease: populations, individuals, and health professionals. *Lancet*. 2019;394:685-696.
22. Munzel T, et al. Environmental Stressors and Cardiovascular Health: Acting Locally for Global Impact in a Changing World: A statement of the European Society of Cardiology, the American College of Cardiology, the American Heart Association, and the World Heart Federation. *J Am Coll Cardiol*. 2026.
23. Rapsomaniki E, et al. Prognostic models for stable coronary artery disease based on electronic health record cohort of 102 023 patients. *Eur Heart J*. 2014;35:844-52.
24. Halonen JJ, et al. Road traffic noise is associated with increased cardiovascular morbidity and mortality and all-cause mortality in London. *Eur Heart J*. 2015;36:2653-61.
25. Claeys MJ, et al. Climate and environmental triggers of acute myocardial infarction. *Eur Heart J*. 2016.
26. Badimon L and al. e. Position paper of the European Society of Cardiology–working group of coronary pathophysiology and microcirculation: obesity and heart disease. *Eur Heart J*. 2017.
27. Madhavan MV and al. e. Coronary Artery Disease in Patients ≥ 80 Years of Age. *J Am Coll Cardiol*. 2018.
28. Baggett TP and al. e. Cardiovascular Disease and Homelessness. *J Am Coll Cardiol*. 2018.
29. Rajagopalan S, et al. Personal-Level Protective Actions Against Particulate Matter Air Pollution Exposure: A Scientific Statement From the American Heart Association. *Circulation*. 2020;142:e411-e431.
30. Kaufman JD, et al. Guidance to Reduce the Cardiovascular Burden of Ambient Air Pollutants: A Policy Statement From the American Heart Association. *Circulation*. 2020;142:e432-e447.
31. Vaduganathan M, et al. The Global Burden of Cardiovascular Diseases and Risk: A Compass for Future Health. *J Am Coll Cardiol*. 2022.
32. Koskinas KC, et al. Obesity and cardiovascular disease: an ESC clinical consensus statement. *Eur Heart J*. 2024;45:4063-4098.

33. Gerds E, et al. Sex differences in arterial hypertension. *Eur Heart J*. 2022.
34. George J and al. e. Cardiovascular Effects of Switching From Tobacco Cigarettes to Electronic Cigarettes. *JACC*. 2019.
35. Yao Z, et al. Differential Associations of Cigar, Pipe, and Smokeless Tobacco Use Versus Combustible Cigarette Use With Subclinical Markers of Inflammation, Thrombosis, and Atherosclerosis: The Cross-Cohort Collaboration-Tobacco Working Group. *Circulation*. 2025;151:993-1005.
36. Vadiveloo MK, et al. Ultraprocessed Foods and Their Association With Cardiometabolic Health: Evidence, Gaps, and Opportunities: A Science Advisory From the American Heart Association. *Circulation*. 2025;152:e245-e263.
37. Guasti L, et al. Ultra-processed foods, lifestyle management, and cardiovascular diseases. *Eur Heart J*. 2026;47:3456-3473.
38. Wang YJ, et al. Dietary Sodium Intake and Risk of Cardiovascular Disease: A Systematic Review and Dose-Response Meta-Analysis. *Nutrients*. 2020;12.
39. Ma Y, et al. 24-Hour Urinary Sodium and Potassium Excretion and Cardiovascular Risk. *N Engl J Med*. 2021.
40. Chieng D and al. e. The impact of coffee subtypes on incident cardiovascular disease, arrhythmias, and mortality: long-term outcomes from the UK Biobank. *Eur J Prev Cardiol*. 2022.
41. Marcus GM, et al. Caffeine and Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation*. 2026;154:e345-e355.
42. Trepanowski JF and Ioannidis JPA. Perspective: Limiting Dependence on Nonrandomized Studies and Improving Randomized Trials in Human Nutrition Research: Why and How. *Adv Nutr*. 2018;9:367-377.
43. Carson JAS, et al. Dietary Cholesterol and Cardiovascular Risk: A Science Advisory From the American Heart Association. *Circulation*. 2019;CIR0000000000000743.
44. Seligman HK, et al. A Systematic Review of "Food Is Medicine" Randomized Controlled Trials for Noncommunicable Disease in the United States: A Scientific Statement From the American Heart Association. *Circulation*. 2025.
45. Avis SR, et al. Coronary artery disease in the absence of traditional risk factors: a call for action. *Eur Heart J*. 2021;42:3822-3824.
46. Ference BA, et al. Low-density lipoproteins cause atherosclerotic cardiovascular disease. 1. Evidence from genetic, epidemiologic, and clinical studies. A consensus statement from the European Atherosclerosis Society Consensus Panel. *Eur Heart J*. 2017.
47. Domanski MJ, et al. Time Course of LDL Cholesterol Exposure and Cardiovascular Disease Event Risk. *J Am Coll Cardiol*. 2020;76:1507-1516.
48. Brunner FJ, et al. Application of non-HDL cholesterol for population-based cardiovascular risk stratification: results from the Multinational Cardiovascular Risk Consortium. *Lancet*. 2019.
49. Sniderman AD, et al. The severe hypercholesterolemia phenotype: clinical diagnosis, management, and emerging therapies. *J Am Coll Cardiol*. 2014;63:1935-47.
50. Jorgensen AB, et al. Loss-of-function mutations in APOC3 and risk of ischemic vascular disease. *N Engl J Med*. 2014;371:32-41.
51. Tg, et al. Loss-of-function mutations in APOC3, triglycerides, and coronary disease. *N Engl J Med*. 2014;371:22-31.
52. Klempfner R, et al. Elevated Triglyceride Level Is Independently Associated With Increased All-Cause Mortality in Patients With Established Coronary Heart Disease: Twenty-Two-Year Follow-Up of the Bezafibrate Infarction Prevention Study and Registry. *Circulation Cardiovascular quality and outcomes*. 2016;9:100-8.
53. Laufs U, et al. Clinical review on triglycerides. *Eur Heart J*. 2020;41:99-109c.
54. Balling M and al. e. Elevated LDL Triglycerides and Atherosclerotic Risk. *JACC*. 2023.
55. Ko DT, et al. High-Density Lipoprotein Cholesterol and Cause-Specific Mortality in Individuals Without Previous Cardiovascular Conditions: The CANHEART Study. *J Am Coll Cardiol*. 2016;68:2073-2083.
56. Silbernagel G, et al. High-density lipoprotein cholesterol, coronary artery disease, and cardiovascular mortality. *Eur Heart J*. 2013;34:3563-71.
57. Madsen CM, et al. Extreme high high-density lipoprotein cholesterol is paradoxically associated with high mortality in men and women: two prospective cohort studies. *Eur Heart J*. 2017.
58. Mortensen MB, et al. Low-Density Lipoprotein Cholesterol Is Predominantly Associated With Atherosclerotic Cardiovascular Disease Events in Patients With Evidence of Coronary Atherosclerosis: The Western Denmark Heart Registry. *Circulation*. 2023;147:1053-1063.
59. Bhatia HS, et al. Independence of Lipoprotein(a) and Low-Density Lipoprotein Cholesterol-Mediated Cardiovascular Risk: A Participant-Level Meta-Analysis. *Circulation*. 2025;151:312-321.
60. Mach F, et al. 2025 Focused Update of the 2019 ESC/EAS Guidelines for the management of dyslipidaemias. *Eur Heart J*. 2025.
61. Bellomo TR, et al. Evaluation of Lipoprotein(a) as a Prognostic Marker of Extracoronary Atherosclerotic Vascular Disease Progression. *Circulation*. 2025;152:585-598.
62. Markus MRP, et al. Low-density lipoprotein cholesterol, lipoprotein(a) and high-sensitivity C-reactive protein are independent predictors of cardiovascular events. *Eur Heart J*. 2025;46:3863-3874.
63. MacDougall DE, et al. Lipoprotein(a) and recurrent atherosclerotic cardiovascular events: the US Family Heart Database. *Eur Heart J*. 2025;46:4762-4775.
64. Berman AN and al. e. Lipoprotein(a) and Major Adverse Cardiovascular Events in Patients With or Without Baseline Atherosclerotic Cardiovascular Disease. *JACC*. 2024.

65. Nordestgaard BG and Langsted A. Lipoprotein(a) and cardiovascular disease. *Lancet*. 2024;404:1255-1264.
66. Johannesen CDL, et al. Excess Apolipoprotein B and Cardiovascular Risk in Women and Men. *J Am Coll Cardiol*. 2024;83:2262-2273.
67. Zheutlin AR, et al. Cumulative exposure to atherogenic lipoprotein particles in young adults and subsequent incident atherosclerotic cardiovascular disease. *Eur Heart J*. 2025;46:4302-4312.
68. Sniderman AD, et al. Discordance among apoB, non-high-density lipoprotein cholesterol, and triglycerides: implications for cardiovascular prevention. *Eur Heart J*. 2024;45:2410-2418.
69. Doi T and al. e. Elevated Remnant Cholesterol Reclassifies Risk of Ischemic Heart Disease and Myocardial Infarction. *JACC*. 2022.
70. Tromp TR, et al. Worldwide experience of homozygous familial hypercholesterolaemia: retrospective cohort study. *Lancet*. 2022;399:719-728.
71. Sniderman A and al. e. Key Questions About Familial Hypercholesterolemia. *JACC*. 2022.
72. Choi D, et al. Familial Hypercholesterolemia: Challenges for a High-Risk Population: JACC Focus Seminar 1/3. *J Am Coll Cardiol*. 2023;81:1621-1632.
73. Hu P, et al. Prevalence of Familial Hypercholesterolemia Among the General Population and Patients With Atherosclerotic Cardiovascular Disease: A Systematic Review and Meta-Analysis. *Circulation*. 2020.
74. Wiegman A, et al. Familial hypercholesterolaemia in children and adolescents: a European Atherosclerosis Society consensus statement. *Eur Heart J*. 2026;47:3324-3346.
75. D'Erasmus L, et al. Clinical Implications of Monogenic Versus Polygenic Hypercholesterolemia: Long-Term Response to Treatment, Coronary Atherosclerosis Burden, and Cardiovascular Events. *Journal of the American Heart Association*. 2021;10:e018932.
76. Gidding SS, et al. The Agenda for Familial Hypercholesterolemia: A Scientific Statement From the American Heart Association. *Circulation*. 2015;132:2167-92.
77. Wiegman A, et al. Familial hypercholesterolaemia in children and adolescents: gaining decades of life by optimizing detection and treatment. *Eur Heart J*. 2015;36:2425-37.
78. Bogsrud MP, et al. LDL-cholesterol in newborns and children with genetically verified familial hypercholesterolaemia: implications for cholesterol-based screening. *Eur Heart J*. 2025;46:5261-5269.
79. Damase TR, et al. Established and Emerging Nucleic Acid Therapies for Familial Hypercholesterolemia. *Circulation*. 2024;150:724-735.
80. Cuchel M, et al. Homozygous familial hypercholesterolaemia: new insights and guidance for clinicians to improve detection and clinical management. A position paper from the Consensus Panel on Familial Hypercholesterolaemia of the European Atherosclerosis Society. *Eur Heart J*. 2014;35:2146-57.
81. Cuchel M, et al. 2023 Update on European Atherosclerosis Society Consensus Statement on Homozygous Familial Hypercholesterolaemia: new treatments and clinical guidance. *Eur Heart J*. 2023;44:2277-2291.
82. Sjouke B, et al. Homozygous autosomal dominant hypercholesterolaemia in the Netherlands: prevalence, genotype-phenotype relationship, and clinical outcome. *Eur Heart J*. 2015;36:560-5.
83. Mach F, et al. 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J*. 2019.
84. Watts GF and al. e. Plozasiran for Managing Persistent Chylomicronemia and Pancreatitis Risk. *NEJM*. 2024.
85. Sturm AC, et al. Clinical Genetic Testing for Familial Hypercholesterolemia: JACC Scientific Expert Panel. *J Am Coll Cardiol*. 2018;72:662-680.
86. Khera AV, et al. Diagnostic Yield and Clinical Utility of Sequencing Familial Hypercholesterolemia Genes in Patients With Severe Hypercholesterolemia. *J Am Coll Cardiol*. 2016;67:2578-89.
87. Amor-Salamanca A and al. e. Genetically Confirmed Familial Hypercholesterolemia in Patients With Acute Coronary Syndrome. *J Am Coll Cardiol*. 2017.
88. Damask A, et al. Patients With High Genome-Wide Polygenic Risk Scores for Coronary Artery Disease May Receive Greater Clinical Benefit From Alirocumab Treatment in the ODYSSEY OUTCOMES Trial. *Circulation*. 2020;141:624-636.
89. Marston NA, et al. Predicting Benefit From Evolocumab Therapy in Patients With Atherosclerotic Disease Using a Genetic Risk Score: Results From the FOURIER Trial. *Circulation*. 2020;141:616-623.
90. Wilcox T and al. e. Diabetic Agents, From Metformin to SGLT2 Inhibitors and GLP1 Receptor Agonists. *JACC*. 2020.
91. Jankowski J, et al. Cardiovascular Disease in Chronic Kidney Disease: Pathophysiological Insights and Therapeutic Options. *Circulation*. 2021;143:1157-1172.
92. Marx-Schutt K, et al. Cardiovascular disease in chronic kidney disease. *Eur Heart J*. 2025;46:2148-2160.
93. Reinert MS, et al. Polycystic Ovarian Syndrome and the Long-Term Risk of Arrhythmias. *Circulation*. 2026;153:493-503.
94. Powell-Wiley TM, et al. Obesity and Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation*. 2021;143:e984-e1010.
95. Ruperez C, et al. Obesity accelerates cardiovascular ageing. *Eur Heart J*. 2025;46:2161-2185.
96. Lv Y, et al. Body mass index, waist circumference, and mortality in subjects older than 80 years: a Mendelian randomization study. *Eur Heart J*. 2024;45:2145-2154.

97. Fedchenko M, et al. Long-term outcomes after myocardial infarction in middle-aged and older patients with congenital heart disease-a nationwide study. *Eur Heart J*. 2021;42:2577-2586.
98. Sternheim D and al. e. Myocardial Bridging: Diagnosis, Functional Assessment, and Management: JACC State-of-the-Art Review. *JACC*. 2021.
99. Rodriguez BC, et al. Myocardial bridging: a practical guide for clinicians. *Eur Heart J*. 2026;47:5266-5278.
100. Messerli FH and Burnier M. Cardiovascular disease and uric acid: is the not-so-innocent bystander becoming a true culprit and does the US black box warning for febuxostat indicate that not all uric acid lowering is beneficial? *Eur Heart J*. 2019;40:1787-1789.
101. Stahl EP, et al. Nonalcoholic Fatty Liver Disease and the Heart: JACC State-of-the-Art Review. *J Am Coll Cardiol*. 2019;73:948-963.
102. Cappola AR, et al. Thyroid and Cardiovascular Disease: Research Agenda for Enhancing Knowledge, Prevention, and Treatment. *Thyroid*. 2019;29:760-777.
103. Navi BB, et al. Risk of Arterial Thromboembolism in Patients With Cancer. *J Am Coll Cardiol*. 2017;70:926-938.
104. Prochaska JH, et al. Chronic venous insufficiency, cardiovascular disease, and mortality: a population study. *Eur Heart J*. 2021;42:4157-4165.
105. Libby P and al. e. Inflammation, Immunity, and Infection in Atherothrombosis. *JACC*. 2018.
106. Sager HB and Koenig W. Immune cell-based cardiovascular risk assessment: spotlight on the neutrophil-lymphocyte ratio. *Eur Heart J*. 2021;42:904-906.
107. Ridker PM, et al. C-reactive protein and cardiovascular risk among women with no standard modifiable risk factors: evaluating the 'SMuRF-less but inflamed'. *Eur Heart J*. 2026;47:306-314.
108. Sipila PN, et al. Severe Infection and Risk of Cardiovascular Disease: A Multicohort Study. *Circulation*. 2023;147:1582-1593.
109. Cainzos-Achirica M and al. e. Inflammatory Bowel Disease and Atherosclerotic Cardiovascular Disease. *JACC*. 2020.
110. Garshick MS and al. e. Cardiovascular Risk in Patients With Psoriasis. *JACC*. 2021.
111. Ntsekhe M and Baker JV. Cardiovascular Disease Among Persons Living With HIV: New Insights Into Pathogenesis and Clinical Manifestations in a Global Context. *Circulation*. 2023;147:83-100.
112. van Nimwegen FA, et al. Cardiovascular disease after Hodgkin lymphoma treatment: 40-year disease risk. *JAMA Intern Med*. 2015;175:1007-17.
113. Corban MT and al. e. Antiphospholipid Syndrome Role of Vascular Endothelial Cells and Implications for Risk Stratification and Targeted Therapeutics. *J Am Coll Cardiol*. 2017;69:2317-30.
114. Mahmoodi BK, et al. Association of Factor V Leiden With Subsequent Atherothrombotic Events: A GENIUS-CHD Study of Individual Participant Data. *Circulation*. 2020;142:546-555.
115. McCrindle BW, et al. Diagnosis, Treatment, and Long-Term Management of Kawasaki Disease: A Scientific Statement for Health Professionals From the American Heart Association. *Circulation*. 2017;135:e927-e999.
116. Dalager-Pedersen M, et al. Risk for myocardial infarction and stroke after community-acquired bacteremia: a 20-year population-based cohort study. *Circulation*. 2014;129:1387-96.
117. Kwong JC and al. e. Acute Myocardial Infarction after Laboratory-Confirmed Influenza Infection. *N Engl J Med*. 2018.
118. Wu VC, et al. Syphilis and cardiovascular risk: a Taiwanese registry. *Eur Heart J*. 2024;45:1512-1520.
119. Clerkin KJ, et al. COVID-19 and Cardiovascular Disease. *Circulation*. 2020;141:1648-1655.
120. Imazio M, et al. Myopericardial complications following COVID-19 disease and vaccination: a clinical consensus statement of the European Society of Cardiology Working Group on Myocardial and Pericardial Diseases. *Eur Heart J*. 2025;46:3328-3338.
121. Lee S, et al. Live zoster vaccination and cardiovascular outcomes: a nationwide, South Korean study. *Eur Heart J*. 2025;46:2991-3002.
122. Michelis KC, et al. Coronary artery manifestations of fibromuscular dysplasia. *J Am Coll Cardiol*. 2014;64:1033-46.
123. Saw J, et al. Angiographic and Intracoronary Manifestations of Coronary Fibromuscular Dysplasia. *Circulation*. 2016;133:1548-59.
124. Havers-Borgersen E, et al. Endometriosis and long-term cardiovascular risk: a nationwide Danish study. *Eur Heart J*. 2024;45:4734-4743.
125. Ryden L, et al. Periodontitis Increases the Risk of a First Myocardial Infarction: A Report From the PAROKRANK Study. *Circulation*. 2016;133:576-83.
126. Yeghiazarians Y, et al. Obstructive Sleep Apnea and Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation*. 2021;144:e56-e67.
127. Gale CR, et al. Mental disorders across the adult life course and future coronary heart disease: evidence for general susceptibility. *Circulation*. 2014;129:186-93.
128. Minissian MB and al. e. Association of Spontaneous Preterm Delivery and Future Maternal Cardiovascular Disease. *Circulation*. 2018.
129. Grandi SM and al. e. Cardiovascular Disease-Related Morbidity and Mortality in Women With a History of Pregnancy Complications. Systematic Review and Meta-Analysis. *Circulation*. 2019.

130. Schmidt M, et al. Cardiovascular safety of non-aspirin non-steroidal anti-inflammatory drugs: review and position paper by the working group for Cardiovascular Pharmacotherapy of the European Society of Cardiology. *Eur Heart J*. 2016;37:1015-23.
131. MacIntyre IM, et al. Regular Acetaminophen Use and Blood Pressure in People With Hypertension: The PATH-BP Trial. *Circulation*. 2022;145:416-423.
132. Zeng C, et al. Sodium-containing acetaminophen and cardiovascular outcomes in individuals with and without hypertension. *Eur Heart J*. 2022;43:1743-1755.
133. Roos-Hesselink JW, et al. Contraception and cardiovascular disease. *Eur Heart J*. 2015;36:1728-34, 1734a-1734b.
134. Bhatia N, et al. Cardiovascular Effects of Androgen Deprivation Therapy for the Treatment of Prostate Cancer: ABCDE Steps to Reduce Cardiovascular Disease in Patients With Prostate Cancer. *Circulation*. 2016;133:537-41.
135. Vigen R, et al. Association of testosterone therapy with mortality, myocardial infarction, and stroke in men with low testosterone levels. *JAMA*. 2013;310:1829-36.
136. Baggish AL, et al. Cardiovascular Toxicity of Illicit Anabolic-Androgenic Steroid Use. *Circulation*. 2017;135:1991-2002.
137. Windfeld-Mathiasen J, et al. Cardiovascular Disease in Anabolic Androgenic Steroid Users. *Circulation*. 2025;151:828-834.
138. Lin ST, et al. Association between antipsychotic use and risk of acute myocardial infarction: a nationwide case-crossover study. *Circulation*. 2014;130:235-43.
139. Winkel P, et al. Clarithromycin for stable coronary heart disease increases all-cause and cardiovascular mortality and cerebrovascular morbidity over 10 years in the CLARICOR randomised, blinded clinical trial. *International journal of cardiology*. 2015;182:459-65.
140. Krantz MJ, et al. Cardiovascular Complications of Opioid Use: JACC State-of-the-Art Review. *J Am Coll Cardiol*. 2021;77:205-223.
141. Havakuk O and al. e. The Cardiovascular Effects of Cocaine. *J Am Coll Cardiol*. 2017.
142. Lelieveld J, et al. Cardiovascular disease burden from ambient air pollution in Europe reassessed using novel hazard ratio functions. *Eur Heart J*. 2019;40:1590-1596.
143. Marfella R, et al. Microplastics and Nanoplastics in Atheromas and Cardiovascular Events. *N Engl J Med*. 2024;390:900-910.
144. Rajagopalan S, et al. Air pollution exposure and cardiometabolic risk. *The lancet Diabetes & endocrinology*. 2024;12:196-208.
145. Prattichizzo F, et al. Micro-nanoplastics and cardiovascular diseases: evidence and perspectives. *Eur Heart J*. 2024;45:4099-4110.
146. Desai MY, et al. Prevention, Diagnosis, and Management of Radiation-Associated Cardiac Disease: JACC Scientific Expert Panel. *J Am Coll Cardiol*. 2019;74:905-927.
147. Canoy D, et al. Age at menarche and risks of coronary heart and other vascular diseases in a large UK cohort. *Circulation*. 2015;131:237-44.
148. Gustad LT, et al. Symptoms of anxiety and depression and risk of acute myocardial infarction: the HUNT 2 study. *Eur Heart J*. 2014;35:1394-403.
149. Mostofsky E, et al. Outbursts of anger as a trigger of acute cardiovascular events: a systematic review and meta-analysis. *Eur Heart J*. 2014;35:1404-10.
150. Smyth A, et al. Physical Activity and Anger or Emotional Upset as Triggers of Acute Myocardial Infarction: The INTERHEART Study. *Circulation*. 2016;134:1059-1067.
151. Moazzami K, et al. Higher Activation of the Rostromedial Prefrontal Cortex during Mental Stress Predicts Major Cardiovascular Disease Events in Individuals with Coronary Artery Disease. *Circulation*. 2020.
152. Pulkki-Raback L, et al. Cumulative effect of psychosocial factors in youth on ideal cardiovascular health in adulthood: the Cardiovascular Risk in Young Finns Study. *Circulation*. 2015;131:245-53.
153. Ai S, et al. Causal associations of short and long sleep durations with 12 cardiovascular diseases: linear and nonlinear Mendelian randomization analyses in UK Biobank. *Eur Heart J*. 2021.
154. Domínguez F and al. e. Association of Sleep Duration and Quality With Subclinical Atherosclerosis. *JACC*. 2019.
155. Crum-Cianflone NF, et al. Impact of combat deployment and posttraumatic stress disorder on newly reported coronary heart disease among US active duty and reserve forces. *Circulation*. 2014;129:1813-20.
156. Alahmad B, et al. Associations Between Extreme Temperatures and Cardiovascular Cause-Specific Mortality: Results From 27 Countries. *Circulation*. 2023;147:35-46.
157. Nelson CP, et al. Genetically determined height and coronary artery disease. *N Engl J Med*. 2015;372:1608-18.
158. Masi S, et al. Rate of telomere shortening and cardiovascular damage: a longitudinal study in the 1946 British Birth Cohort. *Eur Heart J*. 2014;35:3296-303.
159. Delp MD, et al. Apollo Lunar Astronauts Show Higher Cardiovascular Disease Mortality: Possible Deep Space Radiation Effects on the Vascular Endothelium. *Sci Rep*. 2016;6:29901.
160. Zhou A, et al. Non-linear Mendelian randomization analyses support a role for vitamin D deficiency in cardiovascular disease risk. *Eur Heart J*. 2022;43:1731-1739.
161. O'Sullivan JW and al. e. Polygenic Risk Scores for Cardiovascular Disease: A Scientific Statement From the American Heart Association. *Circulation*. 2022.

162. Fischer M, et al. Distinct heritable patterns of angiographic coronary artery disease in families with myocardial infarction. *Circulation*. 2005;111:855-62.
163. Natarajan P, et al. Polygenic Risk Score Identifies Subgroup With Higher Burden of Atherosclerosis and Greater Relative Benefit From Statin Therapy in the Primary Prevention Setting. *Circulation*. 2017;135:2091-2101.
164. Ganesh SK, et al. Genetics and genomics for the prevention and treatment of cardiovascular disease: update: a scientific statement from the American Heart Association. *Circulation*. 2013;128:2813-51.
165. Chan K, et al. Association between the chromosome 9p21 locus and angiographic coronary artery disease burden: a collaborative meta-analysis. *J Am Coll Cardiol*. 2013;61:957-70.
166. Tragante V, et al. The impact of susceptibility loci for coronary artery disease on other vascular domains and recurrence risk. *Eur Heart J*. 2013;34:2896-904.
167. Patel RS, et al. Association of Chromosome 9p21 with Subsequent Coronary Heart Disease Events: A GENIUS-CHD Study of Individual Participant Data. *Circ Genom Precis Med*. 2019.
168. Vasan SK, et al. ABO Blood Group and Risk of Thromboembolic and Arterial Disease: A Study of 1.5 Million Blood Donors. *Circulation*. 2016;133:1449-57; discussion 1457.
169. Charchar FJ, et al. Inheritance of coronary artery disease in men: an analysis of the role of the Y chromosome. *Lancet*. 2012;379:915-22.
170. Myocardial Infarction G, et al. Coding Variation in ANGPTL4, LPL, and SVEP1 and the Risk of Coronary Disease. *N Engl J Med*. 2016;374:1134-44.
171. Dewey FE, et al. Inactivating Variants in ANGPTL4 and Risk of Coronary Artery Disease. *N Engl J Med*. 2016;374:1123-33.
172. Dewey FE, et al. Genetic and Pharmacologic Inactivation of ANGPTL3 and Cardiovascular Disease. *N Engl J Med*. 2017;377:211-221.
173. Graham MJ, et al. Cardiovascular and Metabolic Effects of ANGPTL3 Antisense Oligonucleotides. *N Engl J Med*. 2017;377:222-232.
174. Inouye M and al. e. Genomic Risk Prediction of Coronary Artery Disease in 480,000 Adults. Implications for Primary Prevention. *JACC*. 2018.
175. Sun Y, et al. Association of TSR1 Variants and Spontaneous Coronary Artery Dissection. *J Am Coll Cardiol*. 2019;74:167-176.
176. Xu S and al. e. The novel coronary artery disease risk gene JCAD/KIAA1462 promotes endothelial dysfunction and atherosclerosis. *Eur Heart J*. 2019.
177. Hartiala JA, et al. Genome-wide analysis identifies novel susceptibility loci for myocardial infarction. *Eur Heart J*. 2021;42:919-933.
178. Zeng L, et al. Genetically modulated educational attainment and coronary disease risk. *Eur Heart J*. 2019.
179. Levin MG and Rader DJ. Polygenic Risk Scores and Coronary Artery Disease: Ready for Prime Time? *Circulation*. 2020;141:637-640.
180. Allahverdian S, et al. Contribution of intimal smooth muscle cells to cholesterol accumulation and macrophage-like cells in human atherosclerosis. *Circulation*. 2014;129:1551-9.
181. Libby P. Mechanisms of acute coronary syndromes and their implications for therapy. *N Engl J Med*. 2013;368:2004-13.
182. Murthy VL, et al. Effects of sex on coronary microvascular dysfunction and cardiac outcomes. *Circulation*. 2014;129:2518-27.
183. Boden WE, et al. Myocardial ischaemic syndromes: a new nomenclature to harmonize evolving international clinical practice guidelines. *Eur Heart J*. 2024;45:3701-3706.
184. Rajkumar CA, et al. The Role of the Collateral Circulation in Stable Angina: An Invasive Placebo-Controlled Study. *Circulation*. 2025;152:1541-1551.
185. Aziz A and al. e. Sex-Related Differences in Vasomotor Function in Patients With Angina and Unobstructed Coronary Arteries. *J Am Coll Cardiol*. 2017.
186. Andersson HB, et al. Long-term survival and causes of death in patients with ST-elevation acute coronary syndrome without obstructive coronary artery disease. *Eur Heart J*. 2017.
187. Kunadian V, et al. An EAPCI Expert Consensus Document on Ischaemia with Non-Obstructive Coronary Arteries in Collaboration with European Society of Cardiology Working Group on Coronary Pathophysiology & Microcirculation Endorsed by Coronary Vasomotor Disorders International Study Group. *Eur Heart J*. 2020;41:3504-3520.
188. Lemesle G and al. e. Incident Myocardial Infarction and Very Late Stent Thrombosis in Outpatients With Stable Coronary Artery Disease. *J Am Coll Cardiol*. 2017;69:2149-56.
189. Crea F and Libby P. Acute Coronary Syndromes: The Way Forward From Mechanisms to Precision Treatment. *Circulation*. 2017;136:1155-1166.
190. Falk E, et al. Update on acute coronary syndromes: the pathologists' view. *Eur Heart J*. 2013;34:719-28.
191. Narula J, et al. Histopathologic characteristics of atherosclerotic coronary disease and implications of the findings for the invasive and noninvasive detection of vulnerable plaques. *J Am Coll Cardiol*. 2013;61:1041-51.
192. Jiang S and al. e. Identification of High-Risk Coronary Lesions by 3-Vessel Optical Coherence Tomography. *JACC*. 2023.
193. Borovac JA. Percutaneous coronary intervention for non-obstructive vulnerable plaques. *Lancet*. 2024.
194. Libby P and Pasterkamp G. Requiem for the 'vulnerable plaque'. *Eur Heart J*. 2015;36:2984-7.

195. Oemrawsingh RM, et al. Near-infrared spectroscopy predicts cardiovascular outcome in patients with coronary artery disease. *J Am Coll Cardiol*. 2014;64:2510-8.
196. Katritsis DG, et al. Ruptured versus stable plaques in human coronary arteries. *Coron Artery Dis*. 2011;22:345-51.
197. Meteva D, et al. Acute coronary syndrome with plaque erosion: clinical characteristics, mechanisms, and unmet needs. *Eur Heart J*. 2026;47:1403-1414.
198. Kakizaki R, et al. Silent plaque ruptures in non-obstructive lesions of non-infarct-related arteries: a multimodality, serial intracoronary imaging study. *Eur Heart J*. 2026;47:2814-2826.
199. Tomaniak M, et al. Vulnerable plaques and patients: state-of-the-art. *Eur Heart J*. 2020;41:2997-3004.
200. Michel JB, et al. Intraplaque haemorrhages as the trigger of plaque vulnerability. *Eur Heart J*. 2011;32:1977-85, 1985a, 1985b, 1985c.
201. Mizuno K, et al. Angioscopic evaluation of coronary-artery thrombi in acute coronary syndromes. *N Engl J Med*. 1992;326:287-91.
202. Torii D and al. e. Eruptive Calcified Nodules as a Potential Mechanism of Acute Coronary Thrombosis and Sudden Death. *JACC*. 2021.
203. Arbab-Zadeh A, et al. Acute coronary events. *Circulation*. 2012;125:1147-56.
204. Arbab-Zadeh A and Fuster V. The myth of the "vulnerable plaque": transitioning from a focus on individual lesions to atherosclerotic disease burden for coronary artery disease risk assessment. *J Am Coll Cardiol*. 2015;65:846-55.
205. Silvain J, et al. Composition of coronary thrombus in acute myocardial infarction. *J Am Coll Cardiol*. 2011;57:1359-67.
206. Montecucco F, et al. Pathophysiology of ST-segment elevation myocardial infarction: novel mechanisms and treatments. *Eur Heart J*. 2016;37:1268-83.
207. DeFilippis AP, et al. Assessment and Treatment of Patients With Type 2 Myocardial Infarction and Acute Nonischemic Myocardial Injury. *Circulation*. 2019;140:1661-1678.
208. Fernandez-Jimenez R, et al. Pathophysiology Underlying the Bimodal Edema Phenomenon After Myocardial Ischemia/Reperfusion. *J Am Coll Cardiol*. 2015;66:816-28.
209. Fernández-Jiménez R and al. e. Dynamic Edematous Response of the Human Heart to Myocardial Infarction. *Circulation*. 2017.
210. Sjoblom J, et al. Evolution of left ventricular ejection fraction after acute myocardial infarction: implications for implantable cardioverter-defibrillator eligibility. *Circulation*. 2014;130:743-8.
211. Bularga A, et al. Coronary Artery and Cardiac Disease in Patients With Type 2 Myocardial Infarction: A Prospective Cohort Study. *Circulation*. 2022.
212. Ambrose JA, et al. Angiographic progression of coronary artery disease and the development of myocardial infarction. *J Am Coll Cardiol*. 1988;12:56-62.
213. Little WC, et al. Can coronary angiography predict the site of a subsequent myocardial infarction in patients with mild-to-moderate coronary artery disease? *Circulation*. 1988;78:1157-66.
214. Giroud D, et al. Relation of the site of acute myocardial infarction to the most severe coronary arterial stenosis at prior angiography. *Am J Cardiol*. 1992;69:729-32.
215. Maron DJ, et al. Initial Invasive or Conservative Strategy for Stable Coronary Disease. *N Engl J Med*. 2020;382:1395-1407.
216. Williams MC, et al. Low-Attenuation Noncalcified Plaque on Coronary Computed Tomography Angiography Predicts Myocardial Infarction: Results From the Multicenter SCOT-HEART Trial (Scottish Computed Tomography of the HEART). *Circulation*. 2020;141:1452-1462.
217. Finn AV, et al. Concept of vulnerable/unstable plaque. *Arterioscler Thromb Vasc Biol*. 2010;30:1282-92.
218. Qiao JH and Fishbein MC. The severity of coronary atherosclerosis at sites of plaque rupture with occlusive thrombosis. *J Am Coll Cardiol*. 1991;17:1138-42.
219. Kaski JC, et al. Rapid angiographic progression of coronary artery disease in patients with angina pectoris. The role of complex stenosis morphology. *Circulation*. 1995;92:2058-65.
220. Ojio S, et al. Considerable time from the onset of plaque rupture and/or thrombi until the onset of acute myocardial infarction in humans: coronary angiographic findings within 1 week before the onset of infarction. *Circulation*. 2000;102:2063-9.
221. Maddox TM, et al. Nonobstructive coronary artery disease and risk of myocardial infarction. *JAMA*. 2014;312:1754-63.
222. Stone GW, et al. A prospective natural-history study of coronary atherosclerosis. *N Engl J Med*. 2011;364:226-35.
223. Mancini GB, et al. Angiographic disease progression and residual risk of cardiovascular events while on optimal medical therapy: observations from the COURAGE Trial. *Circulation Cardiovascular interventions*. 2011;4:545-52.
224. Puchner SB, et al. High-risk plaque detected on coronary CT angiography predicts acute coronary syndromes independent of significant stenosis in acute chest pain: results from the ROMICAT-II trial. *J Am Coll Cardiol*. 2014;64:684-92.
225. Williams MC and al. e. Coronary Artery Plaque Characteristics Associated With Adverse Outcomes in the SCOT-HEART Study. *JACC*. 2019.
226. Fuchs A, et al. Subclinical Coronary Atherosclerosis and Risk for Myocardial Infarction in a Danish Cohort : A Prospective Observational Cohort Study. *Ann Intern Med*. 2023.

227. Cheng JM, et al. In vivo detection of high-risk coronary plaques by radiofrequency intravascular ultrasound and cardiovascular outcome: results of the ATHEROREMO-IVUS study. *Eur Heart J*. 2014;35:639-47.
228. Tian J, et al. Prevalence and characteristics of TCFA and degree of coronary artery stenosis: an OCT, IVUS, and angiographic study. *J Am Coll Cardiol*. 2014;64:672-80.
229. Vergallo R and Crea F. Atherosclerotic Plaque Healing. *N Engl J Med*. 2020;383:846-857.
230. Hayes SN, et al. Spontaneous Coronary Artery Dissection: Current State of the Science: A Scientific Statement From the American Heart Association. *Circulation*. 2018.
231. Adlam D, et al. European Society of Cardiology, acute cardiovascular care association, SCAD study group: a position paper on spontaneous coronary artery dissection. *Eur Heart J*. 2018.
232. Hayes SN, et al. Spontaneous Coronary Artery Dissection: JACC State-of-the-Art Review. *J Am Coll Cardiol*. 2020;76:961-984.
233. Kim ESH. Spontaneous Coronary-Artery Dissection. *N Engl J Med*. 2020;383:2358-2370.
234. Saw J, et al. Canadian spontaneous coronary artery dissection cohort study: in-hospital and 30-day outcomes. *Eur Heart J*. 2019.
235. Dang QM, et al. The Australian-New Zealand spontaneous coronary artery dissection cohort study: predictors of major adverse cardiovascular events and recurrence. *Eur Heart J*. 2025;46:2012-2023.
236. Tweet MS, et al. Spontaneous Coronary Artery Dissection Associated With Pregnancy. *J Am Coll Cardiol*. 2017;70:426-435.
237. Vijayaraghavan R, et al. Pregnancy-related spontaneous coronary artery dissection. *Circulation*. 2014;130:1915-20.
238. Havakuk O, et al. Pregnancy and the Risk of Spontaneous Coronary Artery Dissection: An Analysis of 120 Contemporary Cases. *Circulation Cardiovascular interventions*. 2017;10.
239. Crea F, et al. Mechanisms and diagnostic evaluation of persistent or recurrent angina following percutaneous coronary revascularization. *Eur Heart J*. 2019.
240. Bhasin D, et al. Chest Pain With ST-Segment Elevation in a Young Woman: A Broken Heart? *Circulation*. 2021;143:197-201.
241. Sugiishi M and Takatsu F. Cigarette smoking is a major risk factor for coronary spasm. *Circulation*. 1993;87:76-9.
242. Rezkalla SH and Kloner RA. Cocaine-induced acute myocardial infarction. *Clinical medicine & research*. 2007;5:172-6.
243. Alexander KM, et al. A Shocking Development in a Young Male Athlete With Chest Pain. *Circulation*. 2016;133:756-63.
244. Shibata T, et al. Prevalence, Clinical Features, and Prognosis of Acute Myocardial Infarction Attributable to Coronary Artery Embolism. *Circulation*. 2015;132:241-50.
245. Brambatti M, et al. Eosinophilic Myocarditis: Characteristics, Treatment, and Outcomes. *J Am Coll Cardiol*. 2017;70:2363-2375.
246. Agewall S, et al. ESC working group position paper on myocardial infarction with non-obstructive coronary arteries. *Eur Heart J*. 2017;38:143-153.
247. Tamis-Holland JE, et al. Contemporary Diagnosis and Management of Patients With Myocardial Infarction in the Absence of Obstructive Coronary Artery Disease: A Scientific Statement From the American Heart Association. *Circulation*. 2019;139:e891-e908.
248. Reynolds HR, et al. Coronary Optical Coherence Tomography and Cardiac Magnetic Resonance Imaging to Determine Underlying Causes of Myocardial Infarction With Nonobstructive Coronary Arteries in Women. *Circulation*. 2021;143:624-640.
249. Reynolds HR, et al. Multimodality Imaging to Determine Underlying Causes of Myocardial Infarction With Nonobstructive Coronary Arteries in Women and Men. *Circulation*. 2026;153:1606-1622.
250. Dreyer RP, et al. Myocardial infarction with non-obstructive coronary arteries as compared with myocardial infarction and obstructive coronary disease: outcomes in a Medicare population. *Eur Heart J*. 2020;41:870-878.
251. Reynolds HR, et al. Natural History of Patients with Ischemia and No Obstructive Coronary Artery Disease: The CIAO-ISCHEMIA Study. *Circulation*. 2021.
252. McDiarmid AK, et al. Taxonomy of segmental myocardial systolic dysfunction. *Eur Heart J*. 2016.