

CHRONIC HEART FAILURE: THERAPY

Heart Failure with Reduced LVEF

General Measures

Certain measures may prevent or reduce the risk of heart failure (Tables 1 and 2).¹ Regular **aerobic exercise** is beneficial for patients with stable, non-decompensated heart failure, through various mechanisms.² Progressive **rehabilitation** intervention is also useful in patients who were hospitalized for acute decompensated heart failure.³ **Salt restriction** is usually advisable, especially in patients with stage C and D, although a restriction to less than 100 mmol (<1500 mg/day) did not reduce clinical events.⁴ **Potassium** levels in the intervals 2.8–4.1 mmol/L and 4.8–7.4 mmol/L are associated with an increased short-term mortality in chronic heart failure patients, compared with the reference interval of 4.2–4.4 mmol/L.⁵ **Fluid restriction** to 1.5–2 L/day may be considered only in patients with severe symptoms and hyponatremia. **Weight loss** is advisable, especially in patients with coronary artery disease, and weight loss decreases the risk of heart failure, but extreme wasting is an independent risk factor.^{6, 7} **Alcohol** has a direct toxicity on myocardium and at high doses may cause hypertension and arrhythmias. It should be restricted to 2 units a day, and avoided in alcohol-related dilated cardiomyopathy. **Smoking cessation, avoidance of high altitude** (>1500 m), and **vaccination against influenza and pneumococcus** are recommended. **Pregnancy** carries a high risk in patients with heart failure, and any decision should be discussed on an individual basis. **Contraception** with combined hormonal contraceptives is contraindicated due to fluid retention and increased thrombotic risk. Progesteron-only forms are probably safer but still carry a risk of thrombotic risk and osteopenia.⁸ Copper intrauterine devices or transcervical tubal occlusion are relatively safer.⁸ **Sexual activity** is prohibited only in decompensated or advanced heart failure until stabilization.⁹ Patients with **diabetes** should appropriately treated for blood pressure, lipid and glycaemic control, but avoiding hypoglycaemia potentially associated with drug therapy targeting a HbA_{1c}<6.5%.^{10, 6} **Metformin** can be used as long as LV dysfunction is not severe and renal function adequate (creatinine >1.5 mg/dl in men and >1.4 mg/dl in women).¹¹ *Sodium glucose transporter-2 inhibitors* reduce cardiovascular and total mortality in this setting as well as in patients without diabetes (see Drug Therapy). *Thiazolidinediones* may increase the risk of heart failure and should not be used in patients with symptomatic heart failure.¹² Saxagliptin and alogliptin, selective inhibitors of dipeptidyl peptidase 4 (*DPP-4*), have also been shown to increase the rate of hospitalization for heart failure in most trials.¹³ **Non-steroidal anti-inflammatory drugs** (and especially **selective COX-2 inhibitors**) as well as **corticosteroids** and **herbal preparations** (licorice, ginseng, ma huang) cause salt and water retention and should be used with much

caution. In gout attacks **colchicine** is preferred to NSAIDs in symptomatic heart failure patients. **Aspirin** and other indicated antiplatelet agents, when indicated, should be continued in ischemic patients. Although there has been some evidence that aspirin may interfere with prostaglandin synthesis, leading to a reduced effectiveness of ACE inhibition, the clinical significance of this observation is negligible.¹⁴⁻¹⁶ **Tricyclic antidepressants** are contraindicated due to proarrhythmic potential. The value of enhancement of **erythropoiesis** in anemia is controversial. Erythropoiesis stimulation with darbepoetin alfa corrects hemoglobin deficiency in systolic heart failure but fails to improve mortality or hospitalization rates and may increase thromboembolic risk.¹⁷ A recent report by the American College of Physicians recommends against the use of erythropoiesis-stimulating agents in patients with mild to moderate anaemia and congestive heart failure or coronary heart disease.¹⁸ **Iron deficiency** (ferritin <100 ng/mL or 100–300 ng/mL if transferrin saturation <20%) should be corrected with intravenous iron.^{19, 20} However, IV ferric carboxymaltose reduces symptoms and possibly hospitalization but not mortality.²¹⁻²³ The optimal **serum potassium** concentration in both HFrEF and HFpEF appears to be in the range 4.2–5.0 mmol/L.²⁴ In patients with chronic HF, vitamin D deficiency is common, and high-dose **vitamin D3** supplementation is safe, and may be associated with a favorable effect on cardiac function.²⁵ **Omega-3 polyunsaturated fatty acid** supplementation may be useful.²⁶ Influenza vaccination is associated with a reduced risk of hospitalization.²⁷ **Chemotherapy-induced cardiomyopathy**, in general, can occur acutely soon after treatment, within a few months of treatment, or many years later, and may be prevented with an ARB or ACE inhibitor, and most probably beta blockers such as carvedilol (a beta blocker with antioxidant properties).²⁸⁻³¹ Most cardiotoxicity after anthracycline therapy occurs within the first year and is associated with anthracycline dose and LVEF status at the end of treatment. Early detection and prompt drug therapy are important for substantial recovery of cardiac function.³² See Cardio-oncology for a detailed discussion. The Domain Management Framework provides a structured approach to optimizing HF care in older adults with HF. Tools such as the Medication Appropriateness Index and the STOPP/ START criteria can be helpful for prioritizing medications in this patient group.³³

Patients with heart failure should be enrolled in a multidisciplinary-care management program to reduce the risk of heart failure hospitalization.

Drug Therapy

Beta blockers and Renin Angiotensin System Inhibitors (RASi) ie angiotensin converting enzyme inhibitors (ACEI) or angiotensin receptor blockers (ARB) are the cornerstones of heart failure therapy (Figures 1-7, Tables 3-10). Adherence to maximum recommended doses is important.³⁴ Information on these drugs is also provided in the Chapter on Therapy of Hypertension.

For patients with ***Chronic Kidney Disease and T2 Diabetes or albuminuria***, RASi and SGLT2i should be used as first-line therapy. If albuminuria persists, nonsteroidal mineralocorticoid receptor antagonist (MRA) or a GLP-1–based therapy should be added for further kidney and cardiovascular protection.³⁵

Angiotensin converting enzyme inhibitors (ACEI)

ACEI have multiple pleiotropic actions including improved endothelial function, antiproliferative effects on smooth muscle cells, and antithrombotic effects. They decrease mortality in established heart failure (up to 23%) as well as in post-MI patients with reduced LVEF.³⁶ Enalapril, captopril, lisinopril have been mostly studied in CCF trials, and ramipril and trandolapril in post-MI trials, but their benefits appear to be a class effect without differences between tissue or non-tissue ACEI. Patients with heart failure may have increased plasma angiotensin II, and consequently, poor prognosis despite chronic ACE inhibitor therapy.³⁷

Angiotensin receptor blockers (ARB)

ARB bind competitively to the AT1 receptor. ARB also reduce mortality in heart failure, are equally effective with ACEI in randomized trials,³⁸ and very suitable for patients unable to tolerate ACEI due to cough (occurs in 10% in whites, higher in Chinese). Mainly studied agents in CCF are valsartan and candesartan. Irbesartan has been mainly studied in hypertension and diabetic nephropathy trials. Losartan has produced controversial results on mortality at low doses. In patients with LVEF $\leq 40\%$ who remain symptomatic on ACEI and beta blockers, addition of eplerenone is preferred to an ARB. The addition of an ARB (or renin inhibitor) to the combination of an ACE inhibitor and a mineralocorticoid antagonist is not recommended because of the risk of renal dysfunction and hyperkalaemia. In hemodialysis patients with heart failure and LVEF $\leq 40\%$ the addition of telmisartan to ACE significantly reduced all-cause and cardiac mortality, and hospital stays.³⁹ ARBs are not removed by dialysis and the benefits of the combination may be seen even in normotensive patients.³⁹

Contraindications to the use of ACEI/ARB are bilateral renal stenosis, serum potassium > 5 mmol/l, and serum creatinine > 2.5 mg/dl (relative contraindication). A history of angioedema or development of severe cough are contraindications for an ACEI. Angioedema is extremely rare with ARB. With GFR < 30 mL/min/1.73 m², an ARB can still be appropriate, particularly with albuminuric CKD. With eGFR < 15 mL/min/1.73 m², consider reducing/stopping if there are uremic symptoms or other problems.⁴⁰ An increase in serum creatinine of $< 50\%$ above baseline, as long as eGFR remains > 15 mL/min/1.73 m², is considered acceptable.⁴¹ ACEI and ARB are not contraindicated in hospitalized patients with Covid-19.^{42, 43} In adults with CKM syndrome stage 2 to 3 who have CKD and T2D, or CKD without T2D but with UACR ≥ 30 mg/g, and with eGFR ≥ 30

mL/min/1.73 m², use of a RASi (eg, ACEi or ARB) at the maximum tolerated dose is recommended to reduce the loss of kidney function and to lower the risk of CVD.

Angiotensin-neprilysin inhibition (ARNI)

Neprilysin is a neutral endopeptidase responsible for the breakdown of several vasoactive peptides such as natriuretic peptides, bradykinin, and adrenomedullin that can attenuate vasoconstriction and sodium retention, and retard cardiac and vascular hypertrophy and remodelling.⁴⁴ The PARADIGM-HF trial has reported a 15% reduction in total and 20% reduction in cardiovascular mortality over 27 months with LCZ696, a combination of the neprilysin inhibitor **sacubitril** and **valsartan** (200 mg bd which is equivalent to 160 mg of valsartan).⁴⁵ Angiotensin-neprilysin inhibition (ARNI) also appears to prevent the clinical progression of surviving patients with heart failure more effectively than angiotensin-converting enzyme inhibition.⁴⁶ ARNI has also been found superior to enalapril in reducing both sudden cardiac deaths and deaths from worsening heart failure,⁴⁷ as well as causing less hyperkalemia when combined with an aldosterone antagonist.⁴⁸ The combination may also be useful in patients with mildly reduced LVEF.⁴⁹ Moderate eGFR decline that may be seen is not consistently associated with adverse outcomes, and should not deter continuation of sacubitril/valsartan or stall uptitration.⁵⁰

Mineralocorticoid receptor antagonists (MRA)

The secondary hyperaldosteronism seen in patients with heart failure promote sodium retention, and endothelial dysfunction, and may contribute to myocardial fibrosis. Both the selective agent **eplerenone** (25-200 mg od) and the nonselective antagonist **spironolactone** (25-200 mg od) reduce mortality in post-MI patients with LVEF<40% and severe or mild symptoms, or patients with advanced heart failure.^{51, 52} Spironolactone may cause painful gynecomastia in men (antagonist of aldosterone and androgen and progesterone receptors).

Finerenone, a novel non-steroidal MRA, with higher selectivity towards the mineralocorticoid receptor, improved heart failure outcomes in patients with chronic kidney disease and type 2 diabetes, irrespective of a history of HF.⁵³ Renal function and serum potassium levels must be closely monitored due to the risk of hyperkalemia especially when these agents are used together with an ACE or ARB. Correction of hyperkalemia with potassium binding agents such as the nonabsorbable polymer *patiromer*, or the inorganic *sodium zirconium cyclosilicate*, that might allow continuation of MRA and ACEi/ARBs is under study.^{54, 55}

Contraindications to aldosterone antagonists are hyperkalemia (K >5 mmol/l), creatinine>2.5 mg/dl, and concomitant use of a combination of ACE and ARB (relative contraindication).

Aliskiren, a direct renin inhibitor, is not beneficial when added to standard therapy and cannot be used as an alternative to ACEi/ARBs.^{56 57}

Beta blockers

Beta blockers up-regulate β -1 receptor density, blunt norepinephrine and renin production, and mitigate production of deleterious cytokines, including tumor necrosis factor alpha. In addition, β -adrenergic blockade induces effective anti-tumour T cell responses.⁵⁸ They reduce mortality (up to 35%) in heart failure in addition to that offered by ACEI in patients with sinus rhythm. Carvedilol, bisoprolol, and long-acting metoprolol unequivocally reduce mortality.⁵⁹⁻⁶¹ In patients with LVEF $\leq$ 30%, QRS $\geq$ 130 ms, and LBBB, carvedilol has been found more effective than metoprolol in reducing hospitalization and ventricular arrhythmias.⁶² In addition, co-administration of carvedilol with enalapril has been reported to prevent the cardiotoxicity of anthracycline chemotherapy.²⁸ Nebivolol have also been found efficacious in the elderly.⁶³ Xamoterol, a beta blocker with intrinsic sympathomimetic activity and bucindolol, a non-selective beta blocker, have not been found beneficial. Paradoxically, the efficacy of beta blockers has been found reduced in the presence of concomitant AF.⁶⁴ Beta blockers are started on low doses (ie carvedilol 3.125 mg bd) and titrated upwards every 2 weeks. In patients with newly diagnosed heart failure it is safe to use either an ACEI or a beta blocker as first-line therapy.⁶¹

Contraindications to the use of beta blockers are severe asthma, second or third degree AV block and sick sinus syndrome (bradycardia $<$ 50 bpm) without permanent pacing, pheochromocytoma or cocaine use (unopposed α -receptors causing vasoconstriction). COPD is not a contraindication for their use.^{65, 66} In patients with mild to moderate asthma (not in the acute phase) cardioselective beta blockers may be used in minimum doses.⁶⁷

Loop diuretics

These are natriuretic agents by blocking sodium-potassium-chloride cotransporters in the loop of Henle, dilate blood vessels directly, and activate the renin-angiotensin-aldosterone system. Loop diuretics such as frusemide (160 to 320 mg daily) are the only therapy that can acutely produce symptomatic improvement. However, no clinical trial has assessed their effect on mortality in heart failure. They should not be used alone in heart failure since they do not prevent the progression of disease or maintain clinical stability over time. Torsemide (10-20 mg od) and bumetanide (0.5-5 mg od) have higher and more consistent bioavailability without absorption-limited kinetics in comparison with frusemide.⁶⁸ However, in the TRANSFORM-HF RCT, the effects of torsemide and furosemide on patient-reported outcomes were similar regardless of ejection fraction, previous loop diuretic use, and baseline health status.⁶⁹ Subcutaneous frusemide is also now available and may offer better bioavailability.⁷⁰ Renal function and electrolytes should be regularly measured with the use of diuretics due to the risk of hyperkalemia, elevation of BUN, and dilutional hyponatremia. In patients resistant to loop diuretics a thiazide may be added in combination. Chronic use of loop diuretics can lead to hypertrophy of the more distal nephrons, such that even a low dose

of a thiazide-like diuretic had a large diuretic effect and consequently, a downstream effect on electrolytes and kidney function.

Several effects such as the braking phenomenon (acute reduction in diuretic efficacy with repeated loop diuretic dosing), post-diuretic effect (increased sodium retention after the loop diuretic has worn off), rebound sodium retention (increased distal nephron sodium reabsorption due to factors such as distal tubule hypertrophy), increased angiotensin II levels due to persistent volume reduction, and renal adaptation (hypertrophy and hyperfunction of distal tubule cells causing increased local sodium uptake and aldosterone secretion), lead to **diuretic resistance**.⁷¹ In diuretic resistance, bolus doses of 80 mg of furosemide or infusions at a rate of 10-30 mg/h can be used with a goal of treatment 3-5 L urinary volume per day. Worsening renal function, defined as a rise of serum creatinine >0.4 mg/dL, may not portend a poor prognosis when it occurs in patients with effective decongestion.⁶⁸ By blocking distal tubule sodium reabsorption, thiazides or thiazide-like drugs such as metolazone (up to 10 mg daily for 3-5 days) can also antagonize the renal adaptation to chronic loop diuretic therapy and potentially improve diuretic resistance due to rebound sodium retention, even in patients with renal failure.⁷¹ However, there is increased risk of hypokalemia, worsening of renal function, fluid depletion that may require fluid resuscitation, and hyponatremia. Addition of an aldosterone antagonist may reduce hypokalemia and improve natriuresis. In stable patients on optimal therapy, furosemide discontinuation should be considered.⁷²

Acetazolamide (500 mg od) added to loop diuretics may be beneficial, especially in acute decompensated heart failure.⁷³

Sodium–glucose cotransporter-2 inhibitors (SGLT2i) and glucagon-like peptide-1 (GLP-1)

SGLT2i have been found beneficial in patients with,⁷⁴⁻⁷⁶ or without diabetes.⁷⁶⁻⁷⁸ The agents also improve renal function, with few adverse effects.⁷⁹ SGLT2i are now recommended in patients with CCD and heart failure irrespective of diabetes status (Figure 8). Among patients with Type 2 Diabetes (T2D) with CVD or at increased risk for CVD, in addition to lifestyle modification, weight management, and risk factor control, utilization of cardioprotective antihyperglycemic is recommended, including SGLT2, or glucagon-like peptide-1 (GLP-1)–based therapy, or both, to improve cardiovascular and kidney outcomes.³⁵

Digoxin

Digoxin may still have a role in patients with systolic heart failure and concomitant atrial fibrillation. In patients with sinus rhythm digoxin (concentrations of 0.5-0.8 mcg/L) has been shown to offer a reduction of hospital admissions, without any impact on mortality (DIG trial).^{80 81} Digoxin blocks the sodium-potassium (Na-K) ATPase, raising intracellular sodium, and reducing Ca efflux by sodium-calcium exchange. Ca overload of the sarcoplasmic reticulum may trigger arrhythmias. In a recent analysis of the AFFIRM study, digoxin was

associated with a significant increase in all-cause mortality in patients with AF, even in the presence of HF,⁸² but this finding was not verified in another analysis of the same study.⁸³ If used in patients with HF and AF, serum levels should be <1.0 ng/ml. In the DIG trial, low serum digoxin concentrations (0.5 to 0.8 ng per milliliter) were associated with better clinical outcomes than placebo, whereas concentrations above 1.0 ng per milliliter were associated with worse outcomes.⁸⁴ Treatment with low-dose digitoxin has also led to a lower combined risk of death from any cause or hospital admission for worsening heart failure than placebo among patients with heart failure and reduced ejection fraction who received guideline-directed medical therapy.⁸⁵

Other therapies

Hydralazine and nitrates are inferior to ACE inhibitors. However, in African-American heart failure patients (who generally have low plasma renin concentrations and, thus, are theoretically less responsive to RAAS blockade), hydralazine and nitrates are helpful.⁸⁶ Complications are arthralgia and increased antinuclear antibodies (lupus-like syndrome is rare).

The selective cardiac myosin activator **omecantiv mecarbil** is a myotrope that improves left ventricular function without meaningful changes in blood pressure making it potentially attractive for patients with low systolic BP.⁸⁷ It has been shown to lower incidence of a composite of a heart-failure event or death from cardiovascular causes (GALACTIC-HF RCT).⁸⁸ The benefit was modest but statistically significant. Patients who received the drug had a small increase in the cardiac troponin I level, but not an increased incidence of cardiac ischemia or ventricular arrhythmias.

Heart rate that has been identified as a risk factor in chronic heart failure, and **ivabradine**, a selective inhibitor of hyperpolarisation-activated cyclic-nucleotide-gated *funny* current (If) involved in pacemaking-generation and responsiveness of the sinoatrial node, has been found to improve outcome (composite of cardiac death or hospital admission) in patients with heart failure.^{89, 90} However, an increased incidence of AF has been associated with ivabradine,⁹¹ and the addition of ivabradine in patients with heart failure and CAD did not improve outcomes, despite a reduction in admissions for myocardial infarctions and revascularization.⁹² The eligible patients population, mainly inability to take adequate beta blockade, is a small fraction of those with heart failure overall.⁹⁰

In heart failure, endothelial dysfunction and reactive oxygen species reduce nitric oxide bioavailability, resulting in relative deficiency of soluble guanylate cyclase and reduced cyclic GMP generation. **Vericiguat** enhances the cyclic GMP pathway by sensitizing soluble guanylate cyclase to endogenous nitric oxide and has reduced the incidence of death from cardiovascular causes or hospitalization over a 10-month period (VICTORIA trial).⁹³ In the VICTOR RCT, vericiguat led to a significant 17% relative risk reduction for cardiovascular

death and a 16% relative risk reduction for all-cause death. Similarly, 25% and 29% relative risk reductions were seen for sudden cardiac death and death due to HF, respectively (both significant).⁹⁴

Direct sodium removal with a sodium-free osmotic peritoneal solution represents a new potential therapy for nonrenal sodium and fluid removal in edematous disorders such as heart failure.⁹⁵

Initiation of **lipid-lowering therapy** is not recommended in patients with HF in the absence of other indications for their use.⁹⁶ However, the effect of statins in this setting is controversial.⁹⁷

Allopurinol increases ATP conservation and might be beneficial.⁹⁸ However, and in large population-based cohort study allopurinol use was associated with an increased risk of non-fatal CV events and all-cause mortality compared to benzbromarone.⁹⁹

Aspirin use is associated with an increased risk of HF in patients with or without a previous history of CVDs (HOMAGE database),¹⁰⁰ and it should be avoided especially by nonischaemic patients without any other indication.¹⁰¹ Its role in ischemic patients is controversial.

Verapamil and **diltiazem** are negative inotropes and are contraindicated in patients with heart failure. Dihydropyridine calcium channel blockers such as **amlodipine** and **felodipine** are safer but should be still avoided.¹⁰²

The use of **inotropes** and **phosphodiesterase inhibitors** in chronic heart failure is not indicated since it is associated with increased mortality.¹⁰³ The PROMISE study found that oral milrinone increases mortality,¹⁰⁴ whereas oral enoximone does not offer any improvement.¹⁰⁵ Oral levosimendan may improve symptoms but without any effect on hospitalizations or mortality (PERSIST).¹⁰⁶

Oral **magnesium** may be beneficial in patients with hypomagnesemia (<1.7mg/dL).¹⁰⁷

Influenza vaccination may improve outcome in patients with heart failure.¹⁰⁸

Anticoagulation Heart failure is accompanied by a hypercoagulable state, and patients have a 1-3.5% annual risk of stroke that is independent of atrial fibrillation.^{109 110} However, there are no sufficient data to support the use of **warfarin** in patients in normal sinus rhythm without a history of thromboembolic events or echocardiographic evidence of LV thrombus. A reduced risk of ischemic stroke with warfarin used instead of aspirin or aspirin (WARCEF trial) and clopidogrel (WATCH trial) is offset by an increased risk of major hemorrhage.^{14, 15} No differences in mortality were seen although a subanalysis of the WARCEF demonstrated lower event rate (ischemic stroke, intracerebral hemorrhage, or death combined) in patients <60 years old.¹¹¹ **Rivaroxaban** at a dose of 2.5 mg twice daily was also not associated with a significantly lower rate of death, myocardial infarction, or stroke than placebo among patients with worsening chronic heart failure, LVEF<40%, coronary artery disease, and no atrial

fibrillation (COMMANDER HF trial).¹¹² However, rivaroxaban 2.5 mg bd reduced the risk of stroke (a component of the primary end-point) and should be considered following an episode of worsening HF.¹¹³ In the COMPASS trial on patients with a history of mild to moderate heart failure (30%<LVEF<40%) combination rivaroxaban 2.5 mg BID and aspirin compared with aspirin alone demonstrated similar relative risk reduction but higher absolute risk reduction for major adverse cardiovascular events and mortality compared with those without heart failure.¹¹⁴

Device Therapy

Several device-based therapies have been approved or are under trial for the therapy of HF.¹¹⁵

Implantable Cardioverter Defibrillator (ICD)

In certain patients with heart failure an ICD is indicated for either secondary (after VT/VF) or primary prevention (Table 11-see also Chapter on VT). In trials of *primary prevention* in patients with heart failure, ICD reduced mortality in post-MI patients with LVEF<30% (MADIT II)¹¹⁶ and ischemic or non-ischemic cardiomyopathy with LVEF<35% (SCD-HeFT).¹¹⁷ In ischemic patients, ICD should be considered if indications exist at least 40 days after MI (see Chapters on ACS). In patients with non-ischemic heart failure and LVEF<35% results have been controversial, but there is now evidence that an ICD may be favourable (see Chapter 60 for a detailed discussion). In patients with end-stage heart failure (NYHA IV), the risks of multiple shocks and quality of life deterioration must be carefully weighed against the survival benefits. ICD may aggravate heart failure either by right ventricular apical pacing that produces dyssynchrony, or multiple shocks.

Cardiac Resynchronization Therapy (CRT)

Mechanical dyssynchrony refers to nonsynchronous contraction of the wall segments of the left ventricle (intraventricular) or between the left and right ventricles (interventricular). Dyssynchrony impairs systolic function and ventricular filling, increases wall stress, and worsens mitral regurgitation. It is most readily defined by the presence of QRS widening and LBBB configuration on the electrocardiogram and may be visualized on 2-dimensional echocardiography.

Biventricular pacing

Biventricular pacing by atrial-synchronized pacing of the RV and the LV via the coronary sinus to the basal or midventricular LV region, accomplishes reverse remodeling of the LV. This remodeling effect, as well as survival benefit is greater in non-ischemic patients, although clinical improvement with CRT has been shown in both ischemic and non-ischemic heart failure. In patients with NYHA III or IV heart failure, LVEF<35%, and QRS duration >120 ms in the presence of LBBB, CRT reduces morbidity and mortality (up to 36%) even without a defibrillator (CARE-HF trial, and others) (Table 11, and Figures 8-11).¹¹⁸

Approximately 30-35% of patients are ‘nonresponders’, in whom CRT fails to result in benefit despite appropriate indications for therapy.^{119, 120} The presence of RBBB, initial r wave in lead V1, advanced heart failure (NYHA 4), advanced age, and ischemic cardiomyopathy are predictors of poor outcome with CRT, whereas LBBB and QRS>150 ms, female gender and non-ischemic aetiology are predictors of response.^{118, 121, 122} However, patients with intraventricular conduction delay and QRS duration ≥ 150 ms might benefit from CRT even in the absence of LBBB.¹²³ In patients with LVEF $\leq 35\%$, symptoms of heart failure, but a QRS duration < 120 ms (LESSER-EARTH trial),¹²⁴ or even < 130 ms (Echo CRT),¹²⁵ may be detrimental. Women are probably more likely than men to achieve a benefit with QRS 130-149 ms.¹²⁶ CRT-D may also be beneficial in non-LBBB patients with PR interval prolongation and LVEF $< 30\%$, QRS duration ≥ 130 ms, and mild heart failure.¹²⁷ Patients with QRS > 130 ms might also respond to CRT even if LVEF $> 30\%$,¹²⁸ while CRT may be beneficial even in mildly symptomatic patients (NYHA I or II) and a QRS > 120 msec, especially in presence of LBBB morphology.^{129, 130} CRT is also preferable to RV pacing in the presence of LVEF $< 50\%$ and AV block, but the potential of increased LV lead-related complications should be considered (BLOCK-HF trial).¹³¹ The possibility of increased QT interval should be also considered.¹³²

For device programming an AV delay of 120 ms without VV interval are initially chosen.¹³³ A PR ≥ 230 ms is associated with increased rates of hospitalization and AV optimization is warranted.¹³⁴ A short-paced QRS and tall R in V1 are usually seen in effective CRT.¹³³ In patients in sinus rhythm, a DDD pacing mode is preferred but the base rate should allow sensing of intrinsic sinus rhythm as much as possible to avoid unnecessary atrial pacing.¹³⁵ Rate-adaptive pacing should be programmed off until the presence of significant iatrogenic or intrinsic chronotropic incompetence affecting exercise intolerance is proven, bearing in mind that simple age-related rate-adaptive pacing does not improve exercise capacity and may be disadvantageous in some.¹³⁵ The adaptive CRT algorithm withholds right ventricular pacing during periods of intrinsic conduction with an atrioventricular interval of 200 ms or less and delivers left ventricular-only stimulation synchronised with right ventricular activation (synchronised left ventricular stimulation). Although theoretically useful, its clinical value is not established.¹³⁶

Echocardiography techniques for measurements of intra- and inter-ventricular delay do not reliably identify responders to CRT, and the mortality benefit of CRT may occur in the absence of prospective echocardiographic predictors (PROSPECT and SMART-AV).^{137, 138} In addition to the ability to accurately detect mechanical delay, successful CRT is also affected by myocardial scar burden and lead position relative to dyssynchronous segments. Thus, decisions to use CRT should not be based on currently available echocardiographic/Doppler variables. Echocardiography may be used to identify the AV delay yielding optimal LV

filling, and new indices such as the LV global longitudinal strain may be useful.¹³⁹ Device-based AV and VV optimization algorithms have also been proposed.¹⁴⁰

The degree of LV electric delay, assessed by the QLV interval as measured from the onset of the surface (lead II) QRS to the first major peak (positive or negative) of the LV the LV electrogram may predict MR.¹⁴¹ Super-response, defined as an absolute increase in LVEF $\geq$ 10%, or a decrease in ESVI $\geq$ 30%, or a decrease in EDVI $\geq$ -20% is an indication of favourable outcome after RCT.¹⁴²

In summary, biventricular pacing is clearly indicated in:

SR, QRS $\geq$ 150 msec, LBBB, LVEF $\leq$ 35% and NYHA II - IV

It is also reasonable in LVEF $\leq$ 35%, and QRS $\geq$ 130 ms with LBBB, or non-LBBB but QRS $\geq$ 150 ms.

Randomized studies on CRT have been almost exclusively restricted to patients with SR.

Permanent AF is present in 25-30% of CRT candidates, but there is little evidence that CRT is effective in AF (RAFT trial).¹⁴³ In patients with AF who are treated with CRT, AV nodal ablation (modification) has been suggested as a superior strategy compared to rate-slowng drugs.^{144 145} However, in the CAAN-AF RCT, AVN ablation did not reduce all-cause mortality or nonfatal heart failure events compared with medical rate-control therapy in patients with permanent AF and CRT-D. Pre-ablation average heart rate of these patients, however, was low (72 bpm).¹⁴⁶

Technical issues for lead placement are discussed in the relevant Chapter.

CRT with ICD

When CRT is also indicated in HF patients, a combined device with ICD capabilities (CRT-D) should be implanted.

The incremental benefits of combined CRT plus implantable cardioverter-defibrillator (ICD) devices vs CRT-alone devices in patients with LV systolic dysfunction remain uncertain, especially in patients with non-ischaemic cardiomyopathy.¹⁴⁷⁻¹⁵⁰ In patients with LVEF $\leq$ 35%, advanced heart failure (NYHA class III or IV) due to ischemic or nonischemic cardiomyopathies and a QRS interval $\geq$ 120 msec, the presence of ICD capabilities also reduced mortality (COMPANION).¹⁵¹ This effect was seen in non-ischaemic patients,¹⁵² in contrast with the results of another observational study,¹⁵³ and a meta-analysis.¹⁵⁰

Addition of CRT to patients who already need an ICD also reduces mortality. In the RAFT trial in patients with NYHA II or III heart failure, LVEF $\leq$ 30%, and a QRS $\geq$ 120 msec or paced QRS $\geq$ 200 msec, the addition of CRT to ICD improved survival, albeit at a cost of increased implantation-related complications.¹⁵⁴ The survival benefit was sustained over a 14-year follow-up.¹⁵⁵ In the MADIT-CRT, in patients with ischemic (NYHA I or II) or nonischemic (NYHA II) cardiomyopathy, LVEF $\leq$ 30%, and QRS $\geq$ 130 msec with LBBB

morphology, CRT offered a 11% reduction in mortality compared to ICD alone.¹²⁹ The composite end-point of heart failure or death was lower among patients with a CRT-D system as compared with those with an ICD alone, only when restricted to those with at least 90% biventricular pacing.¹⁵⁶ In a real world retrospective cohort study using National Cardiovascular Registry data linked with Medicare claims, patients who were eligible for CRT-D according to established criteria and who received CRT-D had significantly lower risks for death and readmission than those who received ICD therapy alone.¹⁵⁷ Improvements in both survival and heart failure hospitalizations with CRT-D compared to ICD alone are greatest in patients with a QRS ≥ 180 ms with or without LBBB, whereas patients with a QRS 150-179 ms without LBBB may have no improvement in survival with CRT-D.¹⁵⁸ In pacemaker or ICD patients with significant RV pacing burden and LVEF $\leq 35\%$, upgrade to CRT-D compared with ICD therapy also reduced the combined risk of all-cause mortality.¹⁵⁹ No benefit of CRT-D compared with CRT was seen in an analysis of a German health insurance database on patients without an indication for implantation of an implantable cardioverter defibrillator for secondary prevention.¹⁶⁰

Patients with CRT-D who achieve LVEF normalization ($>50\%$) have very low absolute and relative risk of ventricular arrhythmias and a favorable clinical course within 2.2 years of follow-up. Risk of inappropriate ICD therapy is still present and these patients could be considered for downgrade from CRT-D to CRT-P at time of battery-depletion if no ventricular arrhythmias have occurred.¹⁶¹ In a subsequent meta-analysis, patients with improvement of LVEF $\geq 45\%$ and those with primary prevention indication for ICD appear to be at lowest risk.¹⁶²

Conduction system pacing

Specific ***His bundle pacing*** may offer better results to biventricular pacing by means of cardiac resynchronization.¹⁶³ ***Left bundle branch pacing*** -CRT, in particular, has demonstrated greater LVEF improvement than biventricular pacing-CRT,¹⁶⁴⁻¹⁶⁸ and may be associated with a lower incidence of sustained VT/VF and new-onset AF compared with biventricular pacing in heart failure patients.¹⁶⁹ Two recent RCTs comparing biventricular to conduction system pacing provided conflicting results.^{170, 171} The presence of an antero-septal scar influences the efficiency of conduction system pacing.¹⁷² Thus, the strength of evidence for CRT in heart failure is still greater than what is available to support Conduction system pacing.¹³⁰ The HRS recommendations are presented in Figures 14 and 15.

Additional Device Therapies

Cardiac contractility modulation (CCM) is a device-based therapy for heart failure that involves applying relatively high-voltage (≈ 7.5 V), long-duration (≈ 20 milliseconds), biphasic electric signals to the right ventricular septal wall during the absolute myocardial refractory period. CCM signals do not elicit a new contraction, but they induce reverse LV remodeling

and improve LVEF over time. There has been some evidence that CCM provides benefit, particularly those with LVEF between 25% and 45%.¹⁷³

Placement of an **inter-atrial shunt device** reduces PCWP during exercise in patients with class III or ambulatory class IV HF, EF $\geq 40\%$, exercise PCWP ≥ 25 mm Hg, and PCWP-right atrial pressure gradient ≥ 5 mm Hg, reduced PCWP during exercise,¹⁷⁴ but did not reduce the total rate of heart failure events or improve health status.¹⁷⁵

Cardiac remuscularization with biologic ventricular assist tissue (**BioVAT**) derived from allogeneic induced pluripotent stem cells is a new promising modality.¹⁷⁶

In selected patients with MR, trans-catheter edge to edge repair may be indicated (Figure 12).

Heart Failure with Preserved LVEF

Effective therapies of heart failure with preserved systolic function are not established (Table 12 and Figure 13). Therapies addressing specific patient phenotypes may prove to be a more effective approach than the traditional model of applying a uniform treatment.^{177, 178} Blood pressure control, rate/rhythm control in underlying AF, control of pulmonary congestion with diuretic agents, and revascularization when indicated are recommended. Blood pressure control improves diastolic function irrespective of the type of antihypertensive agent used.¹⁷⁹ **ARBs, beta blockers** to control tachycardia and increase the duration of diastole, diuretics in smaller doses than in systolic heart failure, and calcium channel blockers (**diltiazem or verapamil**) are used for symptomatic relief. Despite a lower rate of hospitalizations, no reduction in mortality has been shown with ACEI or ARBs, or beta blockers.^{180, 181} In addition, an unexpected worsening of renal function with ARB therapy has been reported,^{182, 183} and β -blocker withdrawal improved maximal functional capacity in patients with HFpEF and chronotropic incompetence.¹⁸⁴ **Angiotensin/neprilysin inhibition** may be beneficial,^{49, 185} especially in patients with high blood pressure despite optimum therapy.¹⁸⁶ **SGLT2 inhibitors** should be initiated in all HFpEF patients without contraindications.¹⁸⁷ Empagliflozin reduced the combined risk of cardiovascular death or hospitalization for heart failure in patients with heart failure and preserved ejection fraction, regardless of the presence or absence of diabetes (EMPEROR-Preserved RCT).¹⁸⁸ and similar benefits were reported for dapagliflozin (DELIVER RCT).¹⁸⁹ **Tirzepatide**, a long-acting agonist of glucose-dependent insulinotropic polypeptide and glucagon-like peptide-1 receptors, has been shown to produce $\approx 12\%$ to 21% weight loss in patients with obesity. In the SUMMIT RCT, tirzepatide led to a lower risk of a composite of death from cardiovascular causes or worsening heart failure than placebo and improved health status in patients with heart failure with preserved ejection fraction and obesity.^{190, 191} Spironolactone has not been beneficial in patients already taking

ACEI/ARBs and diuretics (TOPCAT RCT).¹⁹² However, **finerinone**, another nonsteroidal mineralocorticoid receptor antagonist, resulted in a significantly lower rate of a composite of total worsening heart failure events and death from cardiovascular causes than placebo in patients with heart failure and mildly reduced or preserved ejection fraction.¹⁹³ Ivabradine has not been found beneficial for improving exercise capacity.¹⁹⁴ Nitrates are not indicated.¹⁹⁵ Phosphodiesterase 9A (PDE9A) inhibition that is not dependent on nitric oxide levels, as occurs with conventional PDE5A inhibitors used in pulmonary hypertension, has been found promising in animals with induced cardiac hypertrophy and heart failure.¹⁹⁶

Comorbidities in Heart Failure

Coronary Artery Disease

Revascularization is thought to improve survival in patients with ischemic cardiomyopathy and proven myocardial viability by perfusion tests, stress echocardiography, or MRI.^{197, 198} Although this was not verified in both the STICH and the smaller Heart trial,^{199, 200} the value of CABG over medical therapy is now generally accepted. In patients with extensive disease and at least 2 of prognostic factors such as presence of 3-vessel CAD, LVEF below the median (27%), and end-systolic volume index (ESVI) above the median (79 ml/m²), CABG did reduce mortality compared to medical therapy alone.²⁰¹ In addition, over a 10-year follow-up of the STICH, total as well as cardiac mortality was significantly lower in the CABG group (STICHES trial).²⁰² In patients with LVEF ≥ 35% indications for CABG are as in IHD. In patients with ischemic cardiomyopathy subjected to CABG, surgical LV reconstruction reduces LV volumes and LV wall stress but offers no benefit in mortality or hospitalizations (STICH Hypothesis 2 trial).²⁰³ Thus, routine LV reconstruction is not recommended. Ventricular reshaping techniques with new devices are under study.²⁰⁴ Recommendations are provided in Table 13 and Figure 14.

Valve Disease

In heart failure due to valve disease such as AS or MS, surgical or interventional correction may restore almost normal heart function. Functional MR is characterized by annular dilatation and leaflet noncoaptation in the setting of anatomically normal papillary muscles, and valve leaflets. Ischemic or infarct-related, functional MR is typically associated with leaflet tethering and displacement, related to abnormal LV wall motion and geometry or papillary muscle dysfunction. The role of MV surgery (replacement or repair) in this setting remains controversial. Detailed discussions are provided in the Chapters on Valve Disease.

Atrial Fibrillation

AF constitutes a strong and independent risk factor for the development of heart failure, and both conditions frequently co-exist, partly because of common risk factors.²⁰⁵ The prevalence

of atrial fibrillation in heart failure is 5% (NYHA I) to 25-50% (NYHA III/IV) and new-onset AF is most probably associated with increased subsequent risk of mortality and morbidity.²⁰⁶ It is very important to detect those patients with fast AF who have developed heart failure due to tachycardia-induced cardiomyopathy. Atrial kick contributes up to 30% of stroke volume, and an analysis of the CHF-STAT trial indicated that patients with heart failure who maintained sinus rhythm on amiodarone had improved survival,²⁰⁷ but in a randomized comparison patients with AF and congestive heart failure, the rhythm control strategy using cardioversions and drugs has not been shown to be superior to rate control.²⁰⁸

Several RCTS have evaluated catheter ablation for HFrEF. Restoration of sinus rhythm with catheter ablation has been shown to result in significant improvements in ventricular function (CAMFAT trial),²⁰⁹ particularly in the absence of ventricular fibrosis on CMR (CAMERA-MRI trial).²¹⁰ Catheter ablation is also superior to amiodarone in achieving freedom from AF at long-term follow-up and reducing unplanned hospitalization and mortality in patients with heart failure and persistent AF (AATAC trial).²¹¹ In the CASTLE-AF trial, catheter ablation was associated with a significantly lower rate of a composite end point of death from any cause or hospitalization for worsening heart failure compared to medical therapy,²¹² although caveats should be considered.²¹³ In the CASTLE HTx trial, among patients with end-stage heart failure, the combination of catheter ablation and medical therapy was associated with a lower likelihood of a composite of death from any cause, implantation of a left ventricular assist device, or urgent heart transplantation than medical therapy alone.²¹⁴ In two meta-analyses, catheter ablation has been associated with all-cause mortality benefit in patients with AF and heart failure with reduced ejection fraction.^{215, 216}

However, in patients with AF and significantly reduced LVEF ($\leq 35\%$) no benefit of catheter ablation has been demonstrated in the AMICA trial.²¹⁷ This was mainly due to the fact that at 1 year, LVEF had increased in ablation patients to a similar extent as in medical therapy patients, despite the fact that at any time during follow-up, ablation-group patients had more often SR and a lower AF burden than medical therapy patients. In the RAFT AF trial on patients with high-burden AF and HF, there was no statistical difference in all-cause mortality or HF events with ablation-based rhythm control versus rate control at an average 3 year follow-up.²¹⁸ There was a nonsignificant trend for improved outcomes such as LVEF and quality of life with ablation-based rhythm control over rate control. The trial was stopped early because of apparent futility at the time of the interim analysis. Among all relevant RCTs, only CASTLE-AF and AATAC showed a reduction in mortality with catheter ablation.²¹³

In patients with AF and heart failure with a predominantly preserved LVEF, the stringency of rate control may have no effect on cardiovascular morbidity and mortality,²¹⁹ but in patients with heart failure and reduced LVEF, there is greater mortality when resting

heart rate was >100 bpm, and the use of β -blockers is associated with improved survival.^{220 208} β -blockers are the cornerstone of therapy in patients with systolic heart failure, and confer a 27% reduction in the incidence of new-onset AF.²²¹ β -blockers with or without digoxin are used for rate control aiming at a heart rate <100 bpm. A non-dihydropyridine calcium channel blocker with digoxin may be considered in patients with heart failure and preserved LVEF. Catheter ablation is also preferable to a rate-control strategy with beta blockers and/or digoxin by means of symptoms and neurohormonal status,²²² and to amiodarone by means of reduced AF burden and mortality (AATAC trial).²¹¹ AV nodal ablation and biventricular pacing reduces hospitalizations and mortality compared to pharmacological rate control (APAF-CRT trial),²²³ but catheter ablation for pulmonary vein isolation has been found superior to AV nodal ablation and biventricular pacing (PABA-CHF trial).²²⁴ His pacing may be preferable to biventricular pacing.²²⁵

In patients with AF and heart failure dronedarone increased mortality,²²⁶ whereas dofetilide prevented AF recurrences without any effect on mortality.²²⁷ Amiodarone is the only agent that is recommended for prevention of atrial arrhythmias recurrence in patients with heart failure and/or reduced LVEF, but its potential toxicity, as well as concerns about the possibility of increased mortality,²²⁸ should be considered.

Anticoagulation is recommended in all patients with heart failure and AF (by definition CHA₂DS₂-VASC ≥ 1). Ischemic patients are treated with antiplatelet agents when indicated. However, there have been no randomized trials examining the impact of aspirin alone versus placebo in stroke reduction in HF,¹⁰⁹ and routine addition of single-antiplatelet therapy to warfarin in patients with heart failure, AF and vascular disease is not indicated.²²⁹ DOAC, as a class, reduce the risk of systemic embolism, as well as major and intracranial bleeding compared to warfarin in patients with AF and heart failure.²³⁰ The value of CRT in heart failure with AF is not proven (see CRT).

Recommendations are provided In Table 14 and Figure 15.

Conduction disturbances

In patients with HFrEF (<40%) regardless of NYHA class who have an indication for ventricular pacing and high-degree AVB, CRT rather than RV pacing is recommended, regardless of NYHA class, in order to reduce morbidity (ESC 2021 GL on pacing and CRT, I-A). This includes patients with AF.²³¹

Ventricular Arrhythmias

In patients not eligible or who have no access to ICD, amiodarone may reduce the risk of sudden cardiac death but not overall mortality,²³² and it may increase noncardiac mortality in patients with NYHA class III heart failure.²³³ Amiodarone decreased mortality in the GESICA trial (60% non-ischemic patients)²³⁴ but had no significant effect in the SCD-HeFT (70%

ischemic patients).¹¹⁷ It is still recommended in patients with an ICD to reduce shocks, or in patients with VT in whom ICD is not an option. Dronedarone is detrimental in heart failure patients.²²⁶ Other antiarrhythmic drugs are contraindicated in heart failure due to proarrhythmic potential, and negative inotropic effects.

Chronic Kidney Disease (CKD)

The presence of CKD is associated with more severe heart failure, and CKD itself is a strong independent risk factor of poor cardiovascular outcome.³⁵ Although an initial decline in eGFR is observed on initiation of several HFrEF drug classes (angiotensin-converting enzyme inhibitors/angiotensin II receptor blockers/mineralocorticoid receptor antagonists/angiotensin receptor blocker neprilysin inhibitors/sodium glucose cotransporter 2 inhibitors), renal function often stabilizes over time, and the drugs maintain their clinical efficacy. A decline in eGFR in the context of a stable or improving clinical condition should not lead to discontinuation of therapy. Recommendations are provided in Figures 16-18 and Table 15.

Cancer

Several **chemotherapeutic agents** increase the risk of heart failure, and during cancer treatment with potential cardiotoxic therapies, LV systolic function can be monitored through echocardiography.²³⁵⁻²³⁹ Variants in cardiomyopathy genes contribute to cancer therapy–induced cardiomyopathy susceptibility among adult and paediatric patients with cancer.²⁴⁰ The relative risk of heart failure with preserved or midrange LVEF (>40%) increases with cardiac radiation exposure during contemporary conformal breast cancer radiotherapy.²⁴¹ In the PRADA RCT, anthracycline-containing adjuvant therapy for early breast cancer was associated with a decline in LVEF during extended follow-up. Candesartan during adjuvant therapy did not prevent reduction in LVEF at 2 years, but was associated with modest reduction in left ventricular end-diastolic volume and preserved global longitudinal strain.²⁴² In a subsequent meta-analysis, beta-blockers or ACEI/ARBs were significantly associated with higher LVEFs in patients treated with trastuzumab. For anthracyclines, a similar trend was found, although nonsignificant.²⁴³ Thus, treatment with an ACE-I and a beta-blocker (preferably carvedilol) should be started in patients on chemotherapy who develop LV systolic dysfunction, defined as 10% or more absolute reduction in LVEF to a value below 50% (Table 15).²³⁵

Other comorbidities

Recommendations for other comorbidities are provided in Table 17, and Figure 19. Influenza vaccination is associated with a reduced risk of all-cause death in patients with HF and influenza and pneumococcal vaccination, as well as COVID-19 vaccination, when available, should be considered in patients with HF.²³⁵

Advanced Heart Failure

Definition and identification of patients with advanced HF are presented in Table 18, and Figures 20 and 21. The INTERMACS classification is presented in Figure 22. Principles of therapy are indicated in Tables 19-21, and Figure 23. It should be noted that inotropic agents have not shown improved survival in patients with HF. However, regardless of their mechanism of action (eg, inhibition of phosphodiesterase, stimulation of adrenergic or dopaminergic receptors, calcium sensitization), parenteral inotropes remain an option to help the subset of patients with HF who are refractory to other therapies and are suffering consequences from end-organ hypoperfusion.¹⁰² Inotrope dependent patients (ie levels 1 to 3 according to the Interagency Registry for Mechanical Assist Devices-INTERMACS classification) who can be maintained on a single inotropic agent can be discharged home either on long-term IV inotropic support or on oral levosimendan but with detrimental or no effect on mortality.

Acute heart failure and cardiogenic shock are discussed in the relevant Chapter.

Mechanical Circulatory Support

Mechanical circulatory support is now a therapeutic option for patients with advanced HF. Left-ventricular assist devices (LVAD) are portable blood pumps that can be used in patients with end-stage heart failure (Tables 22-24). As bridge to recovery in people with potentially reversible forms of heart failure, such as myocarditis or post-partum cardiomyopathy.

- As bridge to transplantation within the next weeks
- As destination therapy for patients unsuitable for transplantation.

The randomized REMATCH trial in patients with end-stage heart failure unsuitable for transplantation detected a significantly better survival at one year in the device group (52%) than in the medical therapy group (52 vs 25% at one year and 23 vs 8% at two years, respectively).²⁴⁴ The FDA has approved three devices, HVAD, HeartMate II, and HeartMate III as destination therapy for patients with end-stage heart failure, but Medtronic discontinued the production of HVAD in 2021.²⁴⁵ Sufficient recovery to allow device explantation that is used either as bridge or destination therapy has been observed in 5% to 24% of patients.²⁴⁶ Continuous flow devices comprise the overwhelming majority of LVAD implantations in the USA.²⁴⁷ Continuous-flow centrifugal pumps with a magnetically levitated rotor (such as HeartMate 3) are designed to allow an artificial pulse and thus enhance hemocompatibility (i.e. bleeding and thrombosis events) and avert pump thrombosis and strokes.²⁴⁸⁻²⁵¹

Bleeding complications, especially gastrointestinal bleeding, due to aggressive anticoagulation, and, later acquired von Willebrand disease, platelet damage, stroke, device infection, and thrombosis of the inflow cannula, pump chamber, or outflow graft and aortic anastomosis are the most common post-operative adverse events (up to 30%).^{252, 253} In

patients undergoing isolated LVAD implantation, $\geq 20\%$ of them experience acute RV failure, which is a leading cause of premature morbidity and mortality.²⁵⁴ Although the long-term efficacy and safety of LVADs as an alternative to heart transplantation or medical therapy remains uncertain, the limited availability of organs and their use as bridge to recovery, especially before terminal stages of the disease, suggest expansion of their use although concerns about cost-effectiveness have been raised.²⁵⁵⁻²⁵⁸

Recommendations on cardiopulmonary resuscitation in patients with mechanical circulatory support have been published by AHA and are presented in the Chapter on VT.²⁵⁹ Recommendations on arrhythmia management in this setting have also been published by AHA.²⁶⁰ The benefits of pulmonary artery catheters are not proven, but they may be useful in patients with persistent normotensive-hypoperfusion preshock (ie, systolic blood pressure >90 mmHg without inotropes or pressors but with clear evidence of worsened end-organ function).²⁶¹

Heart Transplantation

Approximately 5% and 10% of all patients with heart failure have advanced disease, which is associated with a very high mortality and very poor quality of life (Table 25). Heart transplantation is the treatment of choice for patients with advanced heart failure in the absence of contraindications, with an overall median survival of 12.5 years and conditional survival of 14.8 years for those who survive the first year.²⁶² Approximately 10% of adult heart transplant recipients develop a de novo malignancy between years 1 and 5 after transplantation.²⁶³ Listing criteria and evaluation and management of patients undergoing cardiac transplantation have been described in detail by the International Society for Heart and Lung Transplantation.²⁶⁴ Heart transplantation is indicated in hospitalised patients with status I classification (United Network of Organ Sharing), who need intravenous inotropes or a circulatory support device.²⁵⁶ In ambulatory (status II) patients, transplantation does not improve survival; these patients might benefit from long-term LVAD unloading that may allow recovery of myocardial contractility in patients with dilated cardiomyopathy. According to the 2016 International Society for Heart Lung Transplantation listing criteria for heart transplantation, amyloid heart disease, HIV, and hepatitis B and C infections are no longer contraindications for transplantation.²⁶⁵ Indication for permanent pacing are presented in the relevant Chapter.

Non-cardiac Surgery

Recommendations by the ESC are provided in Table 26.

New therapeutic approaches

A number of new pharmacological pathways, device therapies and biologicals are undergoing advanced stages of investigation with potential for clinical utility in the near future.²⁶⁶ Autologous stem cell intracoronary transplantation may also improve LVEF and clinical outcomes in ischaemic heart failure.²⁶⁷⁻²⁶⁹

Rehabilitation and Sports

Recommendations for exercise and sports in patients with heart failure or heart transplantation are provided in Tables 27-30.

Hospital Discharge and Integration of Care and End of Life

Considerations

Recommendations for discharge and follow-up are provided In Tables 31-35 and Figure 24, and end-of-life considerations in Table 36.

Special Populations

Issues of special populations are presented in Tables 37 and 38.

Pregnancy

Recommendations for pregnant women with heart failure are presented in Tables 39-43.

References

1. Piepoli M, et al. Prevention of heart failure. *Eur Heart J*. 2026;47:4375-4395.
2. Florido R, et al. Six-Year Changes in Physical Activity and the Risk of Incident Heart Failure: The Atherosclerosis Risk in Communities (ARIC) Study. *Circulation*. 2018.
3. Kitzman DW, et al. Physical Rehabilitation for Older Patients Hospitalized for Heart Failure. *N Engl J Med*. 2021;385:203-216.
4. Ezekowitz JA, et al. Reduction of dietary sodium to less than 100 mmol in heart failure (SODIUM-HF): an international, open-label, randomised, controlled trial. *Lancet*. 2022;399:1391-1400.
5. Aldahl M and al. e. Associations of serum potassium levels with mortality in chronic heart failure patients. *Eur Heart J*. 2017.
6. Bozkurt B, et al. Contributory Risk and Management of Comorbidities of Hypertension, Obesity, Diabetes Mellitus, Hyperlipidemia, and Metabolic Syndrome in Chronic Heart Failure: A Scientific Statement From the American Heart Association. *Circulation*. 2016;134:e535-e578.
7. Sundstrom J, et al. Weight Loss and Heart Failure: A Nationwide Study of Gastric Bypass Surgery Versus Intensive Lifestyle Treatment. *Circulation*. 2017;135:1577-1585.
8. Sedlak T, et al. Contraception in patients with heart failure. *Circulation*. 2012;126:1396-400.
9. Steinke EE, et al. Sexual counseling for individuals with cardiovascular disease and their partners: a consensus document from the American Heart Association and the ESC Council on Cardiovascular Nursing and Allied Professions (CCNAP). *Circulation*. 2013;128:2075-96.
10. Zoungas S, et al. Severe hypoglycemia and risks of vascular events and death. *N Engl J Med*. 2010;363:1410-8.
11. Inzucchi SE, et al. Management of hyperglycaemia in type 2 diabetes: a patient-centered approach. Position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia*. 2012;55:1577-96.
12. Page RL, 2nd, et al. Drugs That May Cause or Exacerbate Heart Failure: A Scientific Statement From the American Heart Association. *Circulation*. 2016;134:e32-69.
13. Vijayakumar S and al. e. Glucose-Lowering Therapies and Heart Failure in Type 2 Diabetes Mellitus. Mechanistic Links, Clinical Data, and Future Directions. *Circulation*. 2018.
14. Homma S, et al. Warfarin and aspirin in patients with heart failure and sinus rhythm. *N Engl J Med*. 2012;366:1859-69.
15. Massie BM, et al. Randomized trial of warfarin, aspirin, and clopidogrel in patients with chronic heart failure: the Warfarin and Antiplatelet Therapy in Chronic Heart Failure (WATCH) trial. *Circulation*. 2009;119:1616-24.
16. Teerlink JR, et al. Aspirin Does Not Increase Heart Failure Events in Heart Failure Patients: From the WARCEF Trial. *JACC Heart Fail*. 2017;5:603-610.
17. Swedberg K, et al. Treatment of anemia with darbepoetin alfa in systolic heart failure. *N Engl J Med*. 2013;368:1210-9.
18. Qaseem A, et al. Treatment of anemia in patients with heart disease: a clinical practice guideline from the American College of Physicians. *Ann Intern Med*. 2013;159:770-9.
19. Savarese G, et al. Iron deficiency and cardiovascular disease. *Eur Heart J*. 2023;44:14-27.
20. Kalra PR, et al. Intravenous ferric derisomaltose in patients with heart failure and iron deficiency in the UK (IRONMAN): an investigator-initiated, prospective, randomised, open-label, blinded-endpoint trial. *Lancet*. 2023;400:2199-2209.
21. Mentz RJ, et al. Ferric Carboxymaltose in Heart Failure with Iron Deficiency. *N Engl J Med*. 2023.
22. Ponikowski P, et al. Efficacy of ferric carboxymaltose in heart failure with iron deficiency: an individual patient data meta-analysis. *Eur Heart J*. 2023;44:5077-5091.
23. Packer M, et al. Identification of three mechanistic pathways for iron-deficient heart failure. *Eur Heart J*. 2024;45:2281-2293.
24. Ono R, et al. Optimal serum potassium concentrations in heart failure: an individual patient data meta-analysis. *Eur Heart J*. 2026;47:4480-4499.
25. Witte KK, et al. Effects of Vitamin D on Cardiac Function in Patients With Chronic HF: The VINDICATE Study. *J Am Coll Cardiol*. 2016;67:2593-603.
26. Tavazzi L, et al. Effect of n-3 polyunsaturated fatty acids in patients with chronic heart failure (the GISSI-HF trial): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2008;372:1223-30.
27. Mohseni H, et al. Influenza vaccination and risk of hospitalization in patients with heart failure: a self-controlled case series study. *Eur Heart J*. 2016.
28. Bosch X, et al. Enalapril and carvedilol for preventing chemotherapy-induced left ventricular systolic dysfunction in patients with malignant hemopathies: the OVERCOME trial (preventiOn of left Ventricular dysfunction with Enalapril and caRvedilol in patients submitted to intensive ChemOtherapy for the treatment of Malignant hEmopathies). *J Am Coll Cardiol*. 2013;61:2355-62.
29. Gulati G, et al. Prevention of cardiac dysfunction during adjuvant breast cancer therapy (PRADA): a 2 x 2 factorial, randomized, placebo-controlled, double-blind clinical trial of candesartan and metoprolol. *Eur Heart J*. 2016;37:1671-80.
30. Avila MS and al. e. Carvedilol for Prevention of Chemotherapy-Related Cardiotoxicity. The CECCY Trial. *J Am Coll Cardiol*. 2018.

31. Lewinter C, et al. A systematic review and meta-analysis of beta-blockers and renin-angiotensin system inhibitors for preventing left ventricular dysfunction due to anthracyclines or trastuzumab in patients with breast cancer. *Eur Heart J*. 2022;43:2562-2569.
32. Cardinale D, et al. Early detection of anthracycline cardiotoxicity and improvement with heart failure therapy. *Circulation*. 2015;131:1981-8.
33. Lewsey SC, et al. Strategies for Optimizing Heart Failure Care in the Older Adult: A Scientific Statement From the American Heart Association. *Circulation*. 2026.
34. Ouwerkerk W, et al. Determinants and clinical outcome of uptitration of ACE-inhibitors and beta-blockers in patients with heart failure: a prospective European study. *Eur Heart J*. 2017.
35. Ndumele CE, et al. 2026 AHA/ACC/ADA/ASN Guideline for the Prevention, Detection, Evaluation, and Management of Cardiovascular-Kidney-Metabolic Syndrome: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2026;87:e1889-e2007.
36. Garg R and Yusuf S. Overview of randomized trials of angiotensin-converting enzyme inhibitors on mortality and morbidity in patients with heart failure. Collaborative Group on ACE Inhibitor Trials. *JAMA*. 1995;273:1450-6.
37. Roig E, et al. Clinical implications of increased plasma angiotensin II despite ACE inhibitor therapy in patients with congestive heart failure. *Eur Heart J*. 2000;21:53-7.
38. Lee VC, et al. Meta-analysis: angiotensin-receptor blockers in chronic heart failure and high-risk acute myocardial infarction. *Ann Intern Med*. 2004;141:693-704.
39. Cice G, et al. Effects of telmisartan added to Angiotensin-converting enzyme inhibitors on mortality and morbidity in hemodialysis patients with chronic heart failure a double-blind, placebo-controlled trial. *J Am Coll Cardiol*. 2010;56:1701-8.
40. Kidney Disease: Improving Global Outcomes CKDWG. KDIGO 2024 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int*. 2024;105:S117-S314.
41. Kober L, et al. 2026 ESC Guidelines for the management of heart failure. *Eur Heart J*. 2026.
42. Mehra MR, et al. Cardiovascular Disease, Drug Therapy, and Mortality in Covid-19. *N Engl J Med*. 2020.
43. Sama IE, et al. Circulating plasma concentrations of angiotensin-converting enzyme 2 in men and women with heart failure and effects of renin-angiotensin-aldosterone inhibitors. *Eur Heart J*. 2020;41:1810-1817.
44. Bayes-Genis A, et al. A Test in Context: Neprilysin: Function, Inhibition, and Biomarker. *J Am Coll Cardiol*. 2016;68:639-53.
45. McMurray JJ, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med*. 2014;371:993-1004.
46. Packer M, et al. Angiotensin receptor neprilysin inhibition compared with enalapril on the risk of clinical progression in surviving patients with heart failure. *Circulation*. 2015;131:54-61.
47. Desai AS, et al. Effect of the angiotensin-receptor-neprilysin inhibitor LCZ696 compared with enalapril on mode of death in heart failure patients. *Eur Heart J*. 2015;36:1990-7.
48. Desai AS, et al. Reduced Risk of Hyperkalemia During Treatment of Heart Failure With Mineralocorticoid Receptor Antagonists by Use of Sacubitril/Valsartan Compared With Enalapril: A Secondary Analysis of the PARADIGM-HF Trial. *JAMA Cardiol*. 2016.
49. Vaduganathan M, et al. Sacubitril/valsartan in heart failure with mildly reduced or preserved ejection fraction: a pre-specified participant-level pooled analysis of PARAGLIDE-HF and PARAGON-HF. *Eur Heart J*. 2023;44:2982-2993.
50. Chatur S and al. e. Variation in Renal Function Following Transition to Sacubitril/Valsartan in Patients With Heart Failure. *JACC*. 2023.
51. Pitt B, et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *N Engl J Med*. 1999;341:709-17.
52. Zannad F, et al. Eplerenone in patients with systolic heart failure and mild symptoms. *N Engl J Med*. 2011;364:11-21.
53. Filippatos G, et al. Finerenone Reduces Risk of Incident Heart Failure in Patients With Chronic Kidney Disease and Type 2 Diabetes: Analyses From the FIGARO-DKD Trial. *Circulation*. 2022;145:437-447.
54. Sarwar CM, et al. Hyperkalemia in Heart Failure. *J Am Coll Cardiol*. 2016;68:1575-89.
55. Beldhuis IE, et al. Evidence-Based Medical Therapy in Patients With Heart Failure With Reduced Ejection Fraction and Chronic Kidney Disease. *Circulation*. 2022;145:693-712.
56. Gheorghiade M, et al. Effect of aliskiren on postdischarge mortality and heart failure readmissions among patients hospitalized for heart failure: the ASTRONAUT randomized trial. *JAMA*. 2013;309:1125-35.
57. McMurray JJ, et al. Aliskiren, Enalapril, or Aliskiren and Enalapril in Heart Failure. *N Engl J Med*. 2016;374:1521-32.
58. Globig AM, et al. The beta(1)-adrenergic receptor links sympathetic nerves to T cell exhaustion. *Nature*. 2023;622:383-392.
59. Hjalmarson A, et al. Effects of controlled-release metoprolol on total mortality, hospitalizations, and well-being in patients with heart failure: the Metoprolol CR/XL Randomized Intervention Trial in congestive heart failure (MERIT-HF). MERIT-HF Study Group. *JAMA*. 2000;283:1295-302.
60. Packer M, et al. Effect of carvedilol on survival in severe chronic heart failure. *N Engl J Med*. 2001;344:1651-8.

61. Willenheimer R, et al. Effect on survival and hospitalization of initiating treatment for chronic heart failure with bisoprolol followed by enalapril, as compared with the opposite sequence: results of the randomized Cardiac Insufficiency Bisoprolol Study (CIBIS) III. *Circulation*. 2005;112:2426-35.
62. Ruwald MH, et al. Effect of metoprolol versus carvedilol on outcomes in MADIT-CRT (multicenter automatic defibrillator implantation trial with cardiac resynchronization therapy). *J Am Coll Cardiol*. 2013;61:1518-26.
63. Flather MD, et al. Randomized trial to determine the effect of nebivolol on mortality and cardiovascular hospital admission in elderly patients with heart failure (SENIORS). *Eur Heart J*. 2005;26:215-25.
64. Kotecha D, et al. Efficacy of beta blockers in patients with heart failure plus atrial fibrillation: an individual-patient data meta-analysis. *Lancet*. 2014;384:2235-43.
65. Dransfield MT, et al. Metoprolol for the Prevention of Acute Exacerbations of COPD. *N Engl J Med*. 2019;381:2304-2314.
66. Yang YL, et al. Association of beta-blocker use with survival and pulmonary function in patients with chronic obstructive pulmonary and cardiovascular disease: a systematic review and meta-analysis. *Eur Heart J*. 2020;41:4415-4422.
67. Morales DR, et al. Adverse respiratory effect of acute beta-blocker exposure in asthma: a systematic review and meta-analysis of randomized controlled trials. *Chest*. 2014;145:779-86.
68. Ellison DH and al. e. Diuretics in heart failure. *N Engl J Med*. 2018.
69. Greene SJ, et al. Effect of Torsemide Versus Furosemide on Symptoms and Quality of Life Among Patients Hospitalized for Heart Failure: The TRANSFORM-HF Randomized Clinical Trial. *Circulation*. 2023;148:124-134.
70. Review JS-o-t-A, et al. Diuretic Therapy for Patients With Heart Failure. *JACC*. 2020.
71. Jentzer JC, et al. Combination of loop diuretics with thiazide-type diuretics in heart failure. *J Am Coll Cardiol*. 2010;56:1527-34.
72. Rohde LE, et al. Short-term diuretic withdrawal in stable outpatients with mild heart failure and no fluid retention receiving optimal therapy: a double-blind, multicentre, randomized trial. *Eur Heart J*. 2019;40:3605-3612.
73. Martens P, et al. Decongestion With Acetazolamide in Acute Decompensated Heart Failure Across the Spectrum of Left Ventricular Ejection Fraction: A Prespecified Analysis From the ADVOR Trial. *Circulation*. 2023;147:201-211.
74. Kato ET, et al. Effect of Dapagliflozin on Heart Failure and Mortality in Type 2 Diabetes Mellitus. *Circulation*. 2019;139:2528-2536.
75. Brown AJM, et al. A randomized controlled trial of dapagliflozin on left ventricular hypertrophy in people with type two diabetes: the DAPA-LVH trial. *Eur Heart J*. 2020;41:3421-3432.
76. Cosentino F, et al. Efficacy of Ertugliflozin on Heart Failure-Related Events in Patients with Type 2 Diabetes Mellitus and Established Atherosclerotic Cardiovascular Disease: Results of the VERTIS CV Trial. *Circulation*. 2020.
77. McMurray JJV, et al. Dapagliflozin in Patients with Heart Failure and Reduced Ejection Fraction. *N Engl J Med*. 2019;381:1995-2008.
78. Packer M, et al. Cardiovascular and Renal Outcomes with Empagliflozin in Heart Failure. *N Engl J Med*. 2020.
79. Braunwald E. Gliflozins in the Management of Cardiovascular Disease. *N Engl J Med*. 2022;386:2024-2034.
80. Digitalis Investigation G. The effect of digoxin on mortality and morbidity in patients with heart failure. *N Engl J Med*. 1997;336:525-33.
81. Ambrosy AP, et al. The use of digoxin in patients with worsening chronic heart failure: reconsidering an old drug to reduce hospital admissions. *J Am Coll Cardiol*. 2014;63:1823-32.
82. Whitbeck MG, et al. Increased mortality among patients taking digoxin--analysis from the AFFIRM study. *Eur Heart J*. 2013;34:1481-8.
83. Gheorghiade M, et al. Lack of evidence of increased mortality among patients with atrial fibrillation taking digoxin: findings from post hoc propensity-matched analysis of the AFFIRM trial. *Eur Heart J*. 2013;34:1489-97.
84. Rathore SS, et al. Association of serum digoxin concentration and outcomes in patients with heart failure. *JAMA*. 2003;289:871-8.
85. Bavendiek U, et al. Digitoxin in Patients with Heart Failure and Reduced Ejection Fraction. *N Engl J Med*. 2025.
86. Taylor AL, et al. Combination of isosorbide dinitrate and hydralazine in blacks with heart failure. *N Engl J Med*. 2004;351:2049-57.
87. Metra M, et al. Effects of omecamtiv mecarbil in heart failure with reduced ejection fraction according to blood pressure: the GALACTIC-HF trial. *Eur Heart J*. 2022.
88. Teerlink JR, et al. Cardiac Myosin Activation with Omecamtiv Mecarbil in Systolic Heart Failure. *N Engl J Med*. 2020.
89. Bohm M, et al. Heart rate as a risk factor in chronic heart failure (SHIFT): the association between heart rate and outcomes in a randomised placebo-controlled trial. *Lancet*. 2010;376:886-94.
90. Psotka MA and Teerlink JR. Ivabradine: Role in the Chronic Heart Failure Armamentarium. *Circulation*. 2016;133:2066-75.

91. Martin RI, et al. Atrial fibrillation associated with ivabradine treatment: meta-analysis of randomised controlled trials. *Heart*. 2014;100:1506-10.
92. Fox K, et al. Ivabradine for patients with stable coronary artery disease and left-ventricular systolic dysfunction (BEAUTIFUL): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2008;372:807-16.
93. Armstrong PW and al e. Vericiguat in Patients with Heart Failure and Reduced Ejection Fraction. *NEJM*. 2020.
94. Butler J, et al. Vericiguat and mortality in heart failure and reduced ejection fraction: the VICTOR trial. *Eur Heart J*. 2026;47:683-697.
95. Rao VS and al. e. First-in-Human Experience With Peritoneal Direct Sodium Removal Using a Zero-Sodium Solution. *Circulation*. 2020.
96. Mach F, et al. 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *Eur Heart J*. 2019.
97. Divakaran S and Nohria A. Decoding the link between heart failure and incident cancer. *Eur Heart J*. 2021;42:3060-3062.
98. Hirsch GA, et al. Allopurinol acutely increases adenosine triphosphate energy delivery in failing human hearts. *J Am Coll Cardiol*. 2012;59:802-8.
99. Kang EH, et al. Cardiovascular risk associated with allopurinol vs. benzbromarone in patients with gout. *Eur Heart J*. 2021.
100. Mujaj B, et al. Aspirin use is associated with increased risk for incident heart failure: a patient-level pooled analysis. *ESC Heart Fail*. 2021.
101. Madelaire C, et al. Low-Dose Aspirin in Heart Failure Not Complicated by Atrial Fibrillation: A Nationwide Propensity-Matched Study. *JACC Heart Fail*. 2018;6:156-167.
102. Heidenreich PA, et al. 2022 AHA/ACC/HFSA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. *Circulation*. 2022;145:e895-e1032.
103. Toma M and Starling RC. Inotropic therapy for end-stage heart failure patients. *Curr Treat Options Cardiovasc Med*. 2010;12:409-19.
104. Packer M, et al. Effect of oral milrinone on mortality in severe chronic heart failure. The PROMISE Study Research Group. *N Engl J Med*. 1991;325:1468-75.
105. Metra M, et al. Effects of low-dose oral enoximone administration on mortality, morbidity, and exercise capacity in patients with advanced heart failure: the randomized, double-blind, placebo-controlled, parallel group ESSENTIAL trials. *Eur Heart J*. 2009;30:3015-26.
106. Nieminen MS, et al. Oral levosimendan in patients with severe chronic heart failure --the PERSIST study. *Eur J Heart Fail*. 2008;10:1246-54.
107. Yin Y, et al. Oral magnesium and outcomes in US veterans with heart failure. *Eur Heart J*. 2026;47:80-90.
108. Modin D, et al. Influenza Vaccine in Heart Failure. *Circulation*. 2019;139:575-586.
109. Gurbel PA and Tantry US. Antiplatelet and anticoagulant agents in heart failure: current status and future perspectives. *JACC Heart Fail*. 2014;2:1-14.
110. Abdul-Rahim AH, et al. Risk of Stroke in Chronic Heart Failure Patients Without Atrial Fibrillation: Analysis of the Controlled Rosuvastatin in Multinational Trial Heart Failure (CORONA) and the Gruppo Italiano per lo Studio della Sopravvivenza nell'Insufficienza Cardiaca-Heart Failure (GISSI-HF) Trials. *Circulation*. 2015;131:1486-94; discussion 1494.
111. Homma S, et al. Benefit of warfarin compared with aspirin in patients with heart failure in sinus rhythm: a subgroup analysis of WARCEF, a randomized controlled trial. *Circ Heart Fail*. 2013;6:988-97.
112. Zannad F, et al. Rivaroxaban in Patients with Heart Failure, Sinus Rhythm, and Coronary Disease. *N Engl J Med*. 2018;379:1332-1342.
113. Mehra MR, et al. A comprehensive analysis of the effects of rivaroxaban on stroke or transient ischaemic attack in patients with heart failure, coronary artery disease, and sinus rhythm: the COMMANDER HF trial. *Eur Heart J*. 2019;40:3593-3602.
114. Branch KR, et al. Rivaroxaban With or Without Aspirin in Patients With Heart Failure and Chronic Coronary or Peripheral Artery Disease. *Circulation*. 2019;140:529-537.
115. Fudim M and al. e. Device Therapy in Chronic Heart Failure. *JACC*. 2021.
116. Moss AJ, et al. Prophylactic implantation of a defibrillator in patients with myocardial infarction and reduced ejection fraction. *N Engl J Med*. 2002;346:877-83.
117. Bardy GH, et al. Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure. *N Engl J Med*. 2005;352:225-37.
118. Cleland JG, et al. An individual patient meta-analysis of five randomized trials assessing the effects of cardiac resynchronization therapy on morbidity and mortality in patients with symptomatic heart failure. *Eur Heart J*. 2013;34:3547-56.
119. Prinzen FW, et al. Cardiac resynchronization therapy: state-of-the-art of current applications, guidelines, ongoing trials, and areas of controversy. *Circulation*. 2013;128:2407-18.
120. Leyva F, et al. 20 years of cardiac resynchronization therapy. *J Am Coll Cardiol*. 2014;64:1047-58.
121. Sipahi I, et al. Impact of QRS duration on clinical event reduction with cardiac resynchronization therapy: meta-analysis of randomized controlled trials. *Arch Intern Med*. 2011;171:1454-62.
122. Poole JE, et al. QRS Duration or QRS Morphology: What Really Matters in Cardiac Resynchronization Therapy? *J Am Coll Cardiol*. 2016;67:1104-17.

123. Friedman DJ, et al. Cardiac Resynchronization Therapy Improves Outcomes in Patients With Intraventricular Conduction Delay But Not Right Bundle Branch Block: A Patient-Level Meta-Analysis of Randomized Controlled Trials. *Circulation*. 2023;147:812-823.
124. Thibault B, et al. Cardiac resynchronization therapy in patients with heart failure and a QRS complex <120 milliseconds: the Evaluation of Resynchronization Therapy for Heart Failure (LESSER-EARTH) trial. *Circulation*. 2013;127:873-81.
125. Steffel J, et al. The effect of QRS duration on cardiac resynchronization therapy in patients with a narrow QRS complex: a subgroup analysis of the EchoCRT trial. *Eur Heart J*. 2015;36:1983-9.
126. Zusterzeel R, et al. Cardiac resynchronization therapy in women: US Food and Drug Administration meta-analysis of patient-level data. *JAMA Intern Med*. 2014;174:1340-8.
127. Kutyla V, et al. PR interval identifies clinical response in patients with non-left bundle branch block: a Multicenter Automatic Defibrillator Implantation Trial-Cardiac Resynchronization Therapy substudy. *Circ Arrhythm Electrophysiol*. 2014;7:645-51.
128. Kutyla V, et al. The influence of left ventricular ejection fraction on the effectiveness of cardiac resynchronization therapy: MADIT-CRT (Multicenter Automatic Defibrillator Implantation Trial With Cardiac Resynchronization Therapy). *J Am Coll Cardiol*. 2013;61:936-44.
129. Goldenberg I, et al. Survival with cardiac-resynchronization therapy in mild heart failure. *N Engl J Med*. 2014;370:1694-701.
130. Chung MK, et al. 2023 HRS/APHS/LAHS guideline on cardiac physiologic pacing for the avoidance and mitigation of heart failure. *Heart Rhythm*. 2023;20:e17-e91.
131. Curtis AB, et al. Biventricular pacing for atrioventricular block and systolic dysfunction. *N Engl J Med*. 2013;368:1585-93.
132. Viskin S, et al. Arrhythmogenic Effects of Cardiac Memory. *Circulation*. 2022;146:1170-1181.
133. Daubert C, et al. Avoiding non-responders to cardiac resynchronization therapy: a practical guide. *Eur Heart J*. 2016.
134. Friedman DJ, et al. Association Between a Prolonged PR Interval and Outcomes of Cardiac Resynchronization Therapy: A Report From the National Cardiovascular Data Registry. *Circulation*. 2016;134:1617-1628.
135. Mullens W, et al. Optimized implementation of cardiac resynchronization therapy: a call for action for referral and optimization of care. *Europace*. 2021.
136. Wilkoff BL, et al. Adaptive versus conventional cardiac resynchronisation therapy in patients with heart failure (AdaptResponse): a global, prospective, randomised controlled trial. *Lancet*. 2023;402:1147-1157.
137. Chung ES, et al. Results of the Predictors of Response to CRT (PROSPECT) trial. *Circulation*. 2008;117:2608-16.
138. Ellenbogen KA, et al. Primary results from the SmartDelay determined AV optimization: a comparison to other AV delay methods used in cardiac resynchronization therapy (SMART-AV) trial: a randomized trial comparing empirical, echocardiography-guided, and algorithmic atrioventricular delay programming in cardiac resynchronization therapy. *Circulation*. 2010;122:2660-8.
139. Bax JJ and al.. e. Prognostic implications of left ventricular global longitudinal strain in heart failure patients with narrow QRS complex treated with cardiac resynchronization therapy: a subanalysis of the randomized EchoCRT trial. *Eur Heart J*. 2017;38:720-726.
140. Brugada J and al. e. Contractility sensor-guided optimization of cardiac resynchronization therapy: results from the RESPOND-CRT trial. *Eur Heart J*. 2017;38:730-73.
141. Chatterjee NA, et al. Longer Left Ventricular Electric Delay Reduces Mitral Regurgitation After Cardiac Resynchronization Therapy: Mechanistic Insights From the SMART-AV Study (SmartDelay Determined AV Optimization: A Comparison to Other AV Delay Methods Used in Cardiac Resynchronization Therapy). *Circ Arrhythm Electrophysiol*. 2016;9.
142. Steffel J and Ruschitzka F. Superresponse to cardiac resynchronization therapy. *Circulation*. 2014;130:87-90.
143. Healey JS, et al. Cardiac resynchronization therapy in patients with permanent atrial fibrillation: results from the Resynchronization for Ambulatory Heart Failure Trial (RAFT). *Circ Heart Fail*. 2012;5:566-70.
144. Ganesan AN, et al. Role of AV nodal ablation in cardiac resynchronization in patients with coexistent atrial fibrillation and heart failure a systematic review. *J Am Coll Cardiol*. 2012;59:719-26.
145. Gasparini M, et al. Cardiac resynchronization therapy in patients with atrial fibrillation: the CERTIFY study (Cardiac Resynchronization Therapy in Atrial Fibrillation Patients Multinational Registry). *JACC Heart Fail*. 2013;1:500-7.
146. Sanders P, et al. Cardiac resynchronization therapy with or without atrioventricular node ablation in atrial fibrillation: the CAAN-AF trial. *Eur Heart J*. 2026;47:5128-5138.
147. Al-Khatib SM, et al. Defibrillators: Selecting the Right Device for the Right Patient. *Circulation*. 2016;134:1390-1404.
148. Marijon E, et al. Causes-of-death analysis of patients with cardiac resynchronization therapy: an analysis of the CeRTiTuDe cohort study. *Eur Heart J*. 2015;36:2767-76.
149. Packer M. What causes sudden death in patients with chronic heart failure and a reduced ejection fraction? *Eur Heart J*. 2020;41:1757-1763.
150. Patel D, et al. Cardiac Resynchronization Therapy (CRT) with or without Defibrillation in Patients with Non-Ischemic Cardiomyopathy: A Systematic Review and Meta-analysis. *Circ Arrhythm Electrophysiol*. 2021.

151. Bristow MR, et al. Cardiac-resynchronization therapy with or without an implantable defibrillator in advanced chronic heart failure. *N Engl J Med*. 2004;350:2140-50.
152. Doran B, et al. The Addition of a Defibrillator to Resynchronization Therapy Decreases Mortality in Patients With Nonischemic Cardiomyopathy. *JACC Heart Fail*. 2021.
153. Barra S and al. e. Adding Defibrillation Therapy to Cardiac Resynchronization on the Basis of the Myocardial Substrate. *J Am Coll Cardiol*. 69:1669-78.
154. Tang AS, et al. Cardiac-resynchronization therapy for mild-to-moderate heart failure. *N Engl J Med*. 2010;363:2385-95.
155. Sapp JL, et al. Long-Term Outcomes of Resynchronization-Defibrillation for Heart Failure. *N Engl J Med*. 2024;390:212-220.
156. Ruwald AC, et al. The association between biventricular pacing and cardiac resynchronization therapy-defibrillator efficacy when compared with implantable cardioverter defibrillator on outcomes and reverse remodelling. *Eur Heart J*. 2015;36:440-8.
157. Masoudi FA, et al. Comparative effectiveness of cardiac resynchronization therapy with an implantable cardioverter-defibrillator versus defibrillator therapy alone: a cohort study. *Ann Intern Med*. 2014;160:603-11.
158. Sundaram V, et al. Implantable Cardioverter-Defibrillators With Versus Without Resynchronization Therapy in Patients With a QRS Duration >180 ms. *J Am Coll Cardiol*. 2017;69:2026-2036.
159. Merkely B, et al. Upgrade of right ventricular pacing to cardiac resynchronization therapy in heart failure: a randomized trial. *Eur Heart J*. 2023;44:4259-4269.
160. Hadwiger M, et al. Survival of patients undergoing cardiac resynchronization therapy with or without defibrillator: the RESET-CRT project. *Eur Heart J*. 2022;43:2591-2599.
161. Ruwald MH, et al. Left ventricular ejection fraction normalization in cardiac resynchronization therapy and risk of ventricular arrhythmias and clinical outcomes: results from the Multicenter Automatic Defibrillator Implantation Trial With Cardiac Resynchronization Therapy (MADIT-CRT) trial. *Circulation*. 2014;130:2278-86.
162. Chatterjee NA, et al. Reduced appropriate implantable cardioverter-defibrillator therapy after cardiac resynchronization therapy-induced left ventricular function recovery: a meta-analysis and systematic review. *Eur Heart J*. 2015;36:2780-9.
163. Ferreira Felix I, et al. Conduction system pacing versus biventricular pacing in heart failure with reduced ejection fraction: A systematic review and meta-analysis of randomized controlled trials. *Heart Rhythm*. 2024;21:881-889.
164. Diaz J, C. and al e. Left Bundle Branch Area Pacing Versus Biventricular Pacing as Initial Strategy for Cardiac Resynchronization *JACC*. 2023.
165. Vijayaraman p and al e. Comparison of Left Bundle-Branch Area Pacing to Biventricular Pacing in Candidates for Resynchronization Therapy. *JACC*. 2023.
166. Gin J, et al. Improved outcomes of conduction system pacing in heart failure with reduced ejection fraction: A systematic review and meta-analysis. *Heart Rhythm*. 2023;20:1178-1187.
167. Vijayaraman P, et al. His-Purkinje Conduction System Pacing Optimized Trial of Cardiac Resynchronization Therapy vs Biventricular Pacing: HOT-CRT Clinical Trial. *JACC Clin Electrophysiol*. 2023;9:2628-2638.
168. Diaz JC, et al. Left Bundle Branch Pacing vs Left Ventricular Septal Pacing vs Biventricular Pacing for Cardiac Resynchronization Therapy. *JACC Clin Electrophysiol*. 2023.
169. Herweg B, et al. Arrhythmic Risk in Biventricular Pacing Compared With Left Bundle Branch Area Pacing: Results From the I-CLAS Study. *Circulation*. 2024;149:379-390.
170. Chen X and al. e. Long-Term Outcomes of Left Bundle-Branch Pacing vs Biventricular Pacing in Heart Failure: The HeartSync-LBBP Randomized Clinical Trial. *JAMA Cardiol*. 2026.
171. Zimmerman A, et al. Conduction System vs Biventricular Pacing in Heart Failure: The PhysioSync-HF Randomized Clinical Trial. *JAMA Cardiol*. 2026.
172. Vassallo JA, et al. Left ventricular endocardial activation during right ventricular pacing: effect of underlying heart disease. *J Am Coll Cardiol*. 1986;7:1228-33.
173. Borggreffe M and Mann DL. Cardiac Contractility Modulation in 2018. *Circulation*. 2018;138:2738-2740.
174. Feldman T, et al. A Transcatheter InterAtrial Shunt Device for the Treatment of Heart Failure with Preserved Ejection Fraction (REDUCE LAP-HF I): A Phase 2, Randomized, Sham-Controlled Trial. *Circulation*. 2017.
175. Shah SJ, et al. Atrial shunt device for heart failure with preserved and mildly reduced ejection fraction (REDUCE LAP-HF II): a randomised, multicentre, blinded, sham-controlled trial. *Lancet*. 2022;399:1130-1140.
176. Zimmermann WH, et al. Stem-Cell-Derived Biologic Ventricular Assist Tissue in Heart Failure. *N Engl J Med*. 2026;394:1991-2001.
177. Shah SJ, et al. Phenotype-Specific Treatment of Heart Failure With Preserved Ejection Fraction: A Multiorgan Roadmap. *Circulation*. 2016;134:73-90.
178. Lam CSP, et al. Heart failure with preserved ejection fraction: from mechanisms to therapies. *Eur Heart J*. 2018;39:2780-2792.
179. Solomon SD, et al. Effect of angiotensin receptor blockade and antihypertensive drugs on diastolic function in patients with hypertension and diastolic dysfunction: a randomised trial. *Lancet*. 2007;369:2079-87.
180. Hernandez AF, et al. Clinical effectiveness of beta-blockers in heart failure: findings from the OPTIMIZE-HF (Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients with Heart Failure) Registry. *J Am Coll Cardiol*. 2009;53:184-92.

181. Massie BM, et al. Irbesartan in patients with heart failure and preserved ejection fraction. *N Engl J Med*. 2008;359:2456-67.
182. Damman K, et al. Worsening renal function and outcome in heart failure patients with preserved ejection fraction and the impact of angiotensin receptor blocker treatment. *J Am Coll Cardiol*. 2014;64:1106-13.
183. Damman K, et al. Worsening renal function and outcome in heart failure patients with reduced and preserved ejection fraction and the impact of angiotensin receptor blocker treatment: data from the CHARM-study programme. *Eur J Heart Fail*. 2016;18:1508-1517.
184. Palau P and al. e. Effect of β -Blocker Withdrawal on Functional Capacity in Heart Failure and Preserved Ejection Fraction. *JACC*. 2021.
185. Mc Causland FR, et al. Angiotensin-Neprilysin Inhibition and Renal Outcomes in Heart Failure With Preserved Ejection Fraction. *Circulation*. 2020;142:1236-1245.
186. Jackson AM, et al. Sacubitril-valsartan as a treatment for apparent resistant hypertension in patients with heart failure and preserved ejection fraction. *Eur Heart J*. 2021.
187. 2023 ACC Expert Consensus Decision Pathway on Management of Heart Failure With Preserved Ejection Fraction. *JACC*. 2023.
188. Anker SD, et al. Empagliflozin in Heart Failure with a Preserved Ejection Fraction. *N Engl J Med*. 2021.
189. Solomon SD, et al. Dapagliflozin in Heart Failure with Mildly Reduced or Preserved Ejection Fraction. *N Engl J Med*. 2022.
190. Packer M, et al. Tirzepatide for Heart Failure with Preserved Ejection Fraction and Obesity. *N Engl J Med*. 2025;392:427-437.
191. Zile MR, et al. Effects of Tirzepatide on the Clinical Trajectory of Patients With Heart Failure, Preserved Ejection Fraction, and Obesity. *Circulation*. 2025;151:656-668.
192. Pitt B, et al. Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med*. 2014;370:1383-92.
193. Solomon SD and al. e. Finerenone in Heart Failure with Mildly Reduced or Preserved Ejection Fraction. *NEJM*. 2024.
194. Pal N, et al. Effect of Selective Heart Rate Slowing in Heart Failure With Preserved Ejection Fraction. *Circulation*. 2015;132:1719-25.
195. Redfield MM, et al. Isosorbide Mononitrate in Heart Failure with Preserved Ejection Fraction. *N Engl J Med*. 2015;373:2314-24.
196. Lee DI, et al. Phosphodiesterase 9A controls nitric-oxide-independent cGMP and hypertrophic heart disease. *Nature*. 2015;519:472-6.
197. Allman KC, et al. Myocardial viability testing and impact of revascularization on prognosis in patients with coronary artery disease and left ventricular dysfunction: a meta-analysis. *J Am Coll Cardiol*. 2002;39:1151-8.
198. Buckley O and Di Carli M. Predicting benefit from revascularization in patients with ischemic heart failure: imaging of myocardial ischemia and viability. *Circulation*. 2011;123:444-50.
199. Bonow RO, et al. Myocardial viability and survival in ischemic left ventricular dysfunction. *N Engl J Med*. 2011;364:1617-25.
200. Cleland JG, et al. The Heart Failure Revascularisation Trial (HEART). *Eur J Heart Fail*. 2011;13:227-33.
201. Panza JA, et al. Extent of coronary and myocardial disease and benefit from surgical revascularization in ischemic LV dysfunction [Corrected]. *J Am Coll Cardiol*. 2014;64:553-61.
202. Velazquez EJ, et al. Coronary-artery bypass surgery in patients with left ventricular dysfunction. *N Engl J Med*. 2011;364:1607-16.
203. Jones RH, et al. Coronary bypass surgery with or without surgical ventricular reconstruction. *N Engl J Med*. 2009;360:1705-17.
204. Grossi EA, et al. Outcomes of the RESTOR-MV Trial (Randomized Evaluation of a Surgical Treatment for Off-Pump Repair of the Mitral Valve). *J Am Coll Cardiol*. 2010;56:1984-93.
205. Verma A, et al. Treatment of Patients With Atrial Fibrillation and Heart Failure With Reduced Ejection Fraction. *Circulation*. 2017;135:1547-1563.
206. Swedberg K, et al. Prognostic relevance of atrial fibrillation in patients with chronic heart failure on long-term treatment with beta-blockers: results from COMET. *Eur Heart J*. 2005;26:1303-8.
207. Deedwania PC, et al. Spontaneous conversion and maintenance of sinus rhythm by amiodarone in patients with heart failure and atrial fibrillation: observations from the veterans affairs congestive heart failure survival trial of antiarrhythmic therapy (CHF-STAT). The Department of Veterans Affairs CHF-STAT Investigators. *Circulation*. 1998;98:2574-9.
208. Roy D, et al. Rhythm control versus rate control for atrial fibrillation and heart failure. *N Engl J Med*. 2008;358:2667-77.
209. Hunter RJ ea. A randomized controlled trial of catheter ablation versus medical treatment of atrial fibrillation in heart failure (The CAMFAT trial). *Circulation Arrhythm Electrophysiol*. 2014;7:31-38.
210. Prabhu S, et al. Catheter Ablation Versus Medical Rate control in Atrial Fibrillation and Systolic Dysfunction (CAMERA-MRI). *J Am Coll Cardiol*. 2017.
211. Di Biase L, et al. Ablation Versus Amiodarone for Treatment of Persistent Atrial Fibrillation in Patients With Congestive Heart Failure and an Implanted Device: Results From the AATAC Multicenter Randomized Trial. *Circulation*. 2016;133:1637-44.
212. Marrouche NF, et al. Catheter Ablation for Atrial Fibrillation with Heart Failure. *N Engl J Med*. 2018;378:417-427.

213. Reddy YNV, et al. Management of Atrial Fibrillation Across the Spectrum of Heart Failure With Preserved and Reduced Ejection Fraction. *Circulation*. 2022;146:339-357.
214. Sohns C, et al. Catheter Ablation in End-Stage Heart Failure with Atrial Fibrillation. *N Engl J Med*. 2023;389:1380-1389.
215. Asad ZUA and al. e. Catheter Ablation Versus Medical Therapy for Atrial Fibrillation. A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Circ Arrhythm Electrophysiol*. 2019.
216. Chen S, et al. Rhythm control for patients with atrial fibrillation complicated with heart failure in the contemporary era of catheter ablation: a stratified pooled analysis of randomized data. *Eur Heart J*. 2020;41:2863-2873.
217. Kuck KH, et al. Catheter Ablation Versus Best Medical Therapy in Patients With Persistent Atrial Fibrillation and Congestive Heart Failure: The Randomized AMICA Trial. *Circ Arrhythm Electrophysiol*. 2019;12:e007731.
218. Parkash R, et al. Randomized Ablation-Based Rhythm-Control Versus Rate-Control Trial in Patients with Heart Failure and Atrial Fibrillation: Results from the RAFT-AF trial. *Circulation*. 2022.
219. Mulder BA, et al. Lenient vs. strict rate control in patients with atrial fibrillation and heart failure: a post-hoc analysis of the RACE II study. *Eur J Heart Fail*. 2013;15:1311-8.
220. Li SJ, et al. Prognostic Significance of Resting Heart Rate and Use of beta-Blockers in Atrial Fibrillation and Sinus Rhythm in Patients With Heart Failure and Reduced Ejection Fraction: Findings From the Swedish Heart Failure Registry. *Circ Heart Fail*. 2015;8:871-9.
221. Nasr IA, et al. Prevention of atrial fibrillation onset by beta-blocker treatment in heart failure: a meta-analysis. *Eur Heart J*. 2007;28:457-62.
222. Jones DG, et al. A randomized trial to assess catheter ablation versus rate control in the management of persistent atrial fibrillation in heart failure. *J Am Coll Cardiol*. 2013;61:1894-903.
223. Brignole M, et al. AV junction ablation and cardiac resynchronization for patients with permanent atrial fibrillation and narrow QRS: the APAF-CRT mortality trial. *Eur Heart J*. 2021;42:4731-4739.
224. Khan MN, et al. Pulmonary-vein isolation for atrial fibrillation in patients with heart failure. *N Engl J Med*. 2008;359:1778-85.
225. Huang W, et al. His-bundle pacing vs biventricular pacing following atrioventricular nodal ablation in patients with atrial fibrillation and reduced ejection fraction: A multicenter, randomized, crossover study-The ALTERNATIVE-AF trial. *Heart Rhythm*. 2022;19:1948-1955.
226. Kober L, et al. Increased mortality after dronedarone therapy for severe heart failure. *N Engl J Med*. 2008;358:2678-87.
227. Torp-Pedersen C, et al. Dofetilide in patients with congestive heart failure and left ventricular dysfunction. Danish Investigations of Arrhythmia and Mortality on Dofetilide Study Group. *N Engl J Med*. 1999;341:857-65.
228. Freemantle N, et al. Mixed treatment comparison of dronedarone, amiodarone, sotalol, flecainide, and propafenone, for the management of atrial fibrillation. *Europace*. 2011;13:329-45.
229. Lamberts M, et al. Antithrombotic treatment in patients with heart failure and associated atrial fibrillation and vascular disease: a nationwide cohort study. *J Am Coll Cardiol*. 2014;63:2689-98.
230. Savarese G, et al. Efficacy and Safety of Novel Oral Anticoagulants in Patients With Atrial Fibrillation and Heart Failure: A Meta-Analysis. *JACC Heart failure*. 2016;4:870-880.
231. Glikson M, et al. 2021 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy. *Eur Heart J*. 2021.
232. Piccini JP, et al. Amiodarone for the prevention of sudden cardiac death: a meta-analysis of randomized controlled trials. *Eur Heart J*. 2009;30:1245-53.
233. Packer DL, et al. Impact of implantable cardioverter-defibrillator, amiodarone, and placebo on the mode of death in stable patients with heart failure: analysis from the sudden cardiac death in heart failure trial. *Circulation*. 2009;120:2170-6.
234. Doval HC, et al. Randomised trial of low-dose amiodarone in severe congestive heart failure. Grupo de Estudio de la Sobrevida en la Insuficiencia Cardiaca en Argentina (GESICA). *Lancet*. 1994;344:493-8.
235. 2021 ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure. *Eur Heart J*. 2021.
236. Rao VU and al. e. Clinical Approach to Cardiovascular Toxicity of Oral Antineoplastic Agents. *JACC*. 2021.
237. Bonaca MP and al. e. Myocarditis in the Setting of Cancer Therapeutics. Proposed Case Definitions for Emerging Clinical Syndromes in Cardio-Oncology. *Circulation*. 2019.
238. D'Souza M, et al. The risk of cardiac events in patients receiving immune checkpoint inhibitors: a nationwide Danish study. *Eur Heart J*. 2021;42:1621-1631.
239. Herrmann J, et al. Defining cardiovascular toxicities of cancer therapies: an International Cardio-Oncology Society (IC-OS) consensus statement. *Eur Heart J*. 2022;43:280-299.
240. Garcia-Pavia P, et al. Genetic Variants Associated With Cancer Therapy-Induced Cardiomyopathy. *Circulation*. 2019;140:31-41.
241. Saiki H, et al. Risk of Heart Failure With Preserved Ejection Fraction in Older Women After Contemporary Radiotherapy for Breast Cancer. *Circulation*. 2017;135:1388-1396.
242. Heck SL, et al. Prevention of Cardiac Dysfunction During Adjuvant Breast Cancer Therapy (PRADA): Extended Follow-Up of a 2x2 Factorial, Randomized, Placebo-Controlled, Double-Blind Clinical Trial of Candesartan and Metoprolol. *Circulation*. 2021;143:2431-2440.

243. Lewinter C, et al. A systematic review and meta-analysis of beta-blockers and renin-angiotensin system inhibitors for preventing left ventricular dysfunction due to anthracyclines or trastuzumab in patients with breast cancer. *Eur Heart J*. 2021.
244. Rose EA, et al. Long-term use of a left ventricular assist device for end-stage heart failure. *N Engl J Med*. 2001;345:1435-43.
245. Stehlik J and Kirklin JK. The Long and Winding Road to an Effective Left Ventricular Assist Device: The Demise of Medtronic's HVAD. *Circulation*. 2021;144:509-511.
246. Jakovljevic DG and al. e. Left Ventricular Assist Device as a Bridge to Recovery for Patients With Advanced Heart Failure. *J Am Coll Cardiol*. 2017;69:1924-33.
247. Han JJ, et al. Left Ventricular Assist Devices. *Circulation*. 2018;138:2841-2851.
248. Rogers JG, et al. Intrapericardial Left Ventricular Assist Device for Advanced Heart Failure. *N Engl J Med*. 2017;376:451-460.
249. Pinney SP and al. e. Left Ventricular Assist Devices for Lifelong Support. *J Am Coll Cardiol*. 2017;69.
250. Mehra MR, et al. Two-Year Outcomes with a Magnetically Levitated Cardiac Pump in Heart Failure. *N Engl J Med*. 2018;378:1386-1395.
251. Gustafsson F, et al. Six-month outcomes after treatment of advanced heart failure with a full magnetically levitated continuous flow left ventricular assist device: report from the ELEVATE registry. *Eur Heart J*. 2018;39:3454-3460.
252. Stretch R, et al. National trends in the utilization of short-term mechanical circulatory support: incidence, outcomes, and cost analysis. *J Am Coll Cardiol*. 2014;64:1407-15.
253. Scandroglio AM, et al. Diagnosis and Treatment Algorithm for Blood Flow Obstructions in Patients With Left Ventricular Assist Device. *J Am Coll Cardiol*. 2016;67:2758-68.
254. Konstam MA, et al. Evaluation and Management of Right-Sided Heart Failure: A Scientific Statement From the American Heart Association. *Circulation*. 2018.
255. Rihal CS, et al. 2015 SCAI/ACC/HFSA/STS Clinical Expert Consensus Statement on the Use of Percutaneous Mechanical Circulatory Support Devices in Cardiovascular Care: Endorsed by the American Heart Association, the Cardiological Society of India, and Sociedad Latino Americana de Cardiologia Intervencion; Affirmation of Value by the Canadian Association of Interventional Cardiology-Association Canadienne de Cardiologie d'intervention. *J Am Coll Cardiol*. 2015;65:e7-e26.
256. Guglin M and al. e. Evaluation for Heart Transplantation and LVAD Implantation. *JACC*. 2020.
257. Birks EJ, et al. Prospective Multicenter Study of Myocardial Recovery Using Left Ventricular Assist Devices (RESTAGE-HF [Remission from Stage D Heart Failure]): Medium-Term and Primary End Point Results. *Circulation*. 2020;142:2016-2028.
258. Miller LW, et al. Cost of ventricular assist devices: can we afford the progress? *Circulation*. 2013;127:743-8.
259. Peberdy MA, et al. Cardiopulmonary Resuscitation in Adults and Children With Mechanical Circulatory Support: A Scientific Statement From the American Heart Association. *Circulation*. 2017.
260. Gopinathannair R, et al. Device Therapy and Arrhythmia Management in Left Ventricular Assist Device Recipients: A Scientific Statement From the American Heart Association. *Circulation*. 2019;139:e967-e989.
261. Saxena A, et al. Value of Hemodynamic Monitoring in Patients With Cardiogenic Shock Undergoing Mechanical Circulatory Support. *Circulation*. 2020;141:1184-1197.
262. Crespo-Leiro MG, et al. Heart transplantation: focus on donor recovery strategies, left ventricular assist devices, and novel therapies. *Eur Heart J*. 2022;43:2237-2246.
263. Youn J-C and al. e. Temporal Trends of De Novo Malignancy Development After Heart Transplantation. *J Am Coll Cardiol*. 2018.
264. Mehra MR, et al. Listing criteria for heart transplantation: International Society for Heart and Lung Transplantation guidelines for the care of cardiac transplant candidates--2006. *J Heart Lung Transplant*. 2006;25:1024-42.
265. Mehra MR, et al. The 2016 International Society for Heart Lung Transplantation listing criteria for heart transplantation: A 10-year update. *J Heart Lung Transplant*. 2016;35:1-23.
266. Reddy YNV, et al. Novel approaches to the management of chronic systolic heart failure: future directions and unanswered questions. *Eur Heart J*. 2020;41:1764-1774.
267. Fisher SA, et al. Stem cell therapy for chronic ischaemic heart disease and congestive heart failure. *Cochrane Database Syst Rev*. 2014:CD007888.
268. Afzal MR, et al. Adult Bone Marrow Cell Therapy for Ischemic Heart Disease: Evidence and Insights From Randomized Controlled Trials. *Circ Res*. 2015;117:558-75.
269. Patel AN, et al. Ixmyelocel-T for patients with ischaemic heart failure: a prospective randomised double-blind trial. *Lancet*. 2016;387:2412-21.